

REGISTRATION REPORT

Part A

Risk Management

Product code: BM 20 WP NC

Product name: BOUILLIE BORDELAISE NC 20 K

Chemical active substance:

copper, 200 g/kg

Southern Zone & Interzonal

Zonal Rapporteur Member State: France

NATIONAL ASSESSMENT FRANCE

(authorisation renewal)

Professional uses

Applicant: Industrias Químicas del Vallés S.A. (IQV)

Date: 15/07/2025

Table of Contents

1	Details of the application	4
1.1	Application background	4
1.2	Letters of Access	5
1.3	Justification for submission of tests and studies	5
1.4	Data protection claims	5
2	Details of the authorisation decision	6
2.1	Product identity	6
2.2	Conclusion	6
2.3	Substances of concern for national monitoring	6
2.4	Classification and labelling.....	6
2.4.1	Classification and labelling under Regulation (EC) No 1272/2008	6
2.4.2	Standard phrases under Regulation (EU) No 547/2011.....	7
2.4.3	Other phrases (according to Article 65 (3) of the Regulation (EU) No 1107/2009)	7
2.5	Risk management.....	7
2.5.1	Restrictions linked to the PPP	8
2.5.2	Specific restrictions linked to the intended uses	8
2.6	Intended uses (only NATIONAL GAP)	9
3	Background of authorisation decision and risk management	53
3.1	Physical and chemical properties (Part B, Section 2)	53
3.2	Efficacy (Part B, Section 3)	53
3.3	Methods of analysis (Part B, Section 5).....	53
3.3.1	Analytical method for the formulation	53
3.3.2	Analytical methods for residues.....	53
3.4	Mammalian toxicology (Part B, Section 6)	54
3.4.1	Acute toxicity	54
3.4.2	Operator exposure	54
3.4.3	Worker exposure	59
3.4.4	Bystander exposure	63
3.4.5	Resident exposure	63
3.4.1	Combined exposure	64
3.5	Residues and consumer exposure (Part B, Section 7).....	64
	Summary for BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC)	64
3.6	Environmental fate and behaviour (Part B, Section 8)	66
3.7	Ecotoxicology (Part B, Section 9)	67
3.8	Relevance of metabolites (Part B, Section 10)	68
4	Conclusion of the national comparative assessment (Art. 50 of Regulation (EC) No 1107/2009)	68

5	Further information to permit a decision to be made or to support a review of the conditions and restrictions associated with the authorisation.....	69
5.1.1	Post-authorisation monitoring.....	69
5.1.2	Post-authorisation data requirements	69
Appendix 1	Copy of the product authorisation	70
Appendix 2	Copy of the product label	71

PART A

RISK MANAGEMENT

1 Details of the application

The company INDUSTRIAS QUÍMICAS DEL VALLÉS S.A. has requested a marketing authorisation in France for the product BOUILLIE BORDELAISE NC 20 K (formulation code: BM 20 WP NC), containing 200 g/kg copper¹ (in the form of 733.4 g/kg Bordeaux mixture (CAS n° 8011-63-0)) as fungicide and bactericide for professional uses.

Appendix 1 of this document provides a copy of the product authorisation.

Appendix 2 of this document contains a copy of the product label (draft as proposed by the applicant).

1.1 Application background

The present registration report concerns the evaluation of INDUSTRIAS QUÍMICAS DEL VALLÉS S.A.'s application submitted on 01/04/2019 to market BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) in France (product uses described under point 2.3). France acted as a zonal Rapporteur Member State (zRMS) for field uses and interzonal Rapporteur Member State (izRMS) for greenhouse uses for this request and assessed the application submitted for the re-registration of authorisation after the renewal of approval of the active substances copper compounds of this product in France and in other Member States (MSs) of the European Union.

The present application (2019-4167) was evaluated in France by the French Agency for Food, Environmental and Occupational Health & Safety (Anses), according to the Regulation (EC) n° 1107/2009², the implementing regulations, and French regulations. This application was assessed in the context of the zonal procedure for all MSs of the Southern zone for field uses and for all MSs of the European Union for greenhouse uses, taking into account the worst-case uses ("risk envelope approach")³. When risk mitigation measures were necessary, they are adapted to the situation in France.

The data taken into account are those deemed to be valid either at European level (Review Report and EFSA conclusion) or at zonal/national level. The assessment of BOUILLIE BORDELAISE NC 20 K BM 20 WP NC) has been made using endpoints agreed in the EU peer review of copper compounds. It also includes assessment of data and information related to BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) where those data have not been considered in the EU peer review process.

¹ COMMISSION IMPLEMENTING REGULATION (EU) 2018/1981 of 13 December 2018 renewing the approval of the active substances copper compounds, as candidates for substitution, in accordance with Regulation (EC) No 1107/2009 of the European Parliament and of the Council concerning the placing of plant protection products on the market, and amending the Annex to Commission Implementing Regulation (EU) No 540/2011

² REGULATION (EC) No 1107/2009 of the European Parliament and of the Council of 21 October 2009 concerning the placing of plant protection products on the market and repealing Council Directives 79/117/EEC and 91/414/EEC

³ SANCO document "risk envelope approach", European Commission (14 March 2011). [Guidance document on the preparation and submission of dossiers for plant protection products according to the "risk envelope approach"; SANCO/11244/2011 rev. 5](#)

The conclusions of the assessment published by EFSA 2018^{4,5}, as part of the procedure for the renewal of the approval of copper compounds, based on the available information, identify risk for non-target organisms for the representative uses on grapevine, cucurbits and tomatoes, as well as to workers for the grapevine use.

In the framework of MRL review for copper compounds under Article 12 of Regulation (CE) 396/2005, EFSA published a reasoned opinion (EFSA, 2018⁶). Based on an evaluation of the available data MRL have been proposed and a consumer risk assessment has been conducted. Some information required by the regulation has not been transmitted and a chronic risk for the consumers was identified. Therefore the consumer risk assessment is only tentative and some of the proposed MRL still require a decision by risk managers. Exposure reduction measures could also be investigated.

This part A of the RR presents a summary of essential scientific points upon which recommendations are based and is not intended to show the assessment in detail. The risk assessment conclusions provided in this document are based on the information, data and assessments provided in the Registration Report, Part B Sections 1-10 and Part C, and where appropriate the addendum for France.

The conclusions on the acceptability of risk are based on the criteria provided in Regulation (EU) No 546/2011⁷, and are expressed as “acceptable” or “not acceptable” in accordance with those criteria.

This document also describes the specific conditions of use and labelling required for France for the registration of BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC).

1.2 Letters of Access

The applicant is the owner of data which support the renewal of approval of the active substance. The applicant has provided a letter of access for active substance data. This letter of access is available upon request.

1.3 Justification for submission of tests and studies

According to the applicant: *“The new studies afforded in this dossier, not previously assessed, provided in this dossier are necessary according to the requirements and guidelines into force which didn’t exist when the product was assessed by Uniform Principles in the past.”*

1.4 Data protection claims

2 Where protection for data is being claimed for information supporting registration of BOUILLIE BORDELAISE NC 20 K, it is indicated in the reference lists in Appendix 1 of the Registration Report, Part B Sections 1-7. » ou par « There is no new data

⁴ Peer review of the pesticide risk assessment of the active substance copper compounds Copper(I), copper(II) variants namely copper hydroxide, copper oxychloride, tribasic copper sulfate, copper(I) oxide, Bordeaux mixture, EFSA Journal 2018;16(1):515

⁵ Outcome of the consultation with Member States, the applicant and EFSA on the pesticide risk assessment for copper compounds copper(I), copper(II) variants namely copper hydroxide, copper oxychloride, tribasic copper sulfate, copper(I) oxide, Bordeaux mixture in light of confirmatory data. EFSA supporting publication 2018:EN-1486.

⁶ REASONED OPINION ADOPTED: 1 March 2018. Review of the existing maximum residue levels for copper compounds according to Article 12 of Regulation (EC) No 396/2005 European Food Safety Authority (EFSA).

⁷ COMMISSION REGULATION (EU) No 546/2011 of 10 June 2011 implementing Regulation (EC) No 1107/2009 of the European Parliament and of the Council as regards uniform principles for evaluation and authorisation of plant protection products

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

submitted with this application. Details of the authorisation decision

2.1 Product identity

Product code	BM 20 WP NC
Product name in MS	BOUILLIE BORDELAISE NC 20 K
Authorisation number	<i>N/A : marketing authorisation withdrawn</i>
Kind of use	Professional use
Low risk product (article 47)	No
Function	Fungicide and bactericide
Applicant	INDUSTRIAS QUÍMICAS DEL VALLÉS S.A.
Active substance(s) (incl. content)	copper, 200 g/kg – Bordeaux mixture 733.4 g/kg
Formulation type	Wettable powder [WP]
Coformulants of concern for national authorisations	-
Restrictions related to identity	-
Mandatory tank mixtures	None
Recommended tank mixtures	None

2.2 Conclusion

The evaluation of the application for BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) resulted in the decision to **withdraw** the authorisation.

2.3 Substances of concern for national monitoring

Refer to 5.1.1.

2.4 Classification and labelling

2.4.1 Classification and labelling under Regulation (EC) No 1272/2008

The following classification is proposed in accordance with Regulation (EC) No 1272/2008:

Hazard class(es), categories:	Acute toxicity (inhalation), category 4 Hazardous to the aquatic environment - Acute Hazard, category 1 Hazardous to the aquatic environment - Chronic Hazard, category 1
-------------------------------	---

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Hazard pictograms:	  GHS07 GHS09
Signal word:	Warning
Hazard statement(s):	H332: Harmful if inhaled. H400: Very toxic to aquatic life. H410: Very toxic to aquatic life with long-lasting effects.
Precautionary statement(s):	<i>For the P phrases, refer to the existing legislation</i>
Additional labelling phrases:	-

See Part C for justifications of the classification and labelling proposals.

2.4.2 Standard phrases under Regulation (EU) No 547/2011

N/A : marketing authorisation withdrawn

2.4.3 Other phrases (according to Article 65 (3) of the Regulation (EU) No 1107/2009)

None.

2.5 Risk management

According to the French law and procedures, specific conditions of use are set out in the Decision letter. The French Order of 4 May 2017⁸ provides that:

- unless otherwise stated in the product authorisation, the pre harvest interval (PHI) is at least 3 days;
- unless otherwise stated in the product authorisation, the minimum buffer zone alongside a water body is 5 metres for products applied through spraying or dusting;
- unless otherwise stated in the product authorisation, the minimum re-entry period is 6 hours for field uses and 8 hours for indoor uses.

Drift reduction measures such as low-drift nozzles are not considered within the decision-making process in France. However, non-spraying buffer zones may be reduced under some circumstances as explained in appendix 3 of the above-mentioned French Order.

Finally, the French Order of 26 March 2014⁹ provides that:

- an authorisation granted for a “reference” crop applies also for “related” crops, unless formally stated in the Decision
- the “reference” and “related” crops are defined in Appendix 1 of that French Order.

Thus, at French national level, possible extrapolation of submitted data and the corresponding assessment from “reference” crops to “related” ones are undertaken even if not clearly requested by the applicant in

⁸ Arrêté du 4 mai 2017 relatif à la mise sur le marché et à l'utilisation des produits phytopharmaceutiques et de leurs adjuvants visés à l'article L. 253-1 du code rural et de la pêche maritime, modifié par l'arrêté du 27 décembre 2019 <https://www.legifrance.gouv.fr/eli/arrete/2017/5/4/AGRG1632554A/jo/texte> ; <https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000039686039&categorieLien=id>

⁹ <http://www.legifrance.gouv.fr/eli/arrete/2014/3/26/AGRG1407093A/jo>

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

their dRR, and a conclusion is also reached on the acceptability of the intended uses on those “related” crops. The aim of this Order, mainly based on the EU document on residue data extrapolation¹⁰ is to supply “minor” crops with registered plant protection products.

Therefore the GAP table (Section 2.3) and Decision may include uses on crops not originally requested by the applicant.

Finally, the French Order of 20 November 2021¹¹ on the protection of bees and other pollinating insects and the preservation of pollination services when using plant protection products provides that unless otherwise stated in the product authorisation, use on attractive crops when in flower and on foraging area is forbidden. Specific conditions of application on flowering crops should be respected. As consequences specific SPe 8 may include reference to this order.

The Decision, as reproduced in Appendix 1, takes also into account national provisions, including national mitigation measures.

2.5.1 Restrictions linked to the PPP

The authorisation of the PPP is linked to the following conditions:

Operator protection:	
-	Refer to the Decision in Appendix 1 for the details.
Worker protection:	
-	Refer to the Decision in Appendix 1 for the details.
Integrated pest management (IPM)/sustainable use:	
	-
Environmental protection	
SPe 2	To protect aquatic organisms, do not discharge waste water from soil-less greenhouses directly into surface water.
Precautionary statement for permanent greenhouse	For applications under permanent greenhouse: “May affect pollinators. Avoid unnecessary exposure”.
Other specific restrictions	
Re-entry period	6 hours outdoor 8 hours greenhouse.

2.5.2 Specific restrictions linked to the intended uses

Some of the authorised uses are linked to the following conditions in addition to those listed under point 2.5.1 (mandatory labelling):

None.

¹⁰ SANCO document “guidance document:- Guidelines on comparability, extrapolation, group tolerances and data requirements for setting MRLs”: SANCO/ 7525/VI/95 - rev.9

¹¹ Arrêté du 20 novembre 2021 relatif à la protection des abeilles et des autres insectes pollinisateurs et à la préservation des services de pollinisation lors de l'utilisation des produits phytopharmaceutiques - Légifrance (legifrance.gouv.fr)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

2.6 Intended uses (only NATIONAL GAP)

Please note: The GAP Table below reports the intended uses proposed by the applicant, and possible extrapolation according to French Order of 26 March 2014 (highlighted in green), evaluated and concluded as safe uses by France as zRMS (field uses) and izRMS (greenhouse uses). Those uses are then granted in France.

When the conclusion is “not acceptable””, the intended use is highlighted in grey and the main reason(s) reported in the remarks.

When a use is “acceptable” with GAP restrictions, the modifications of the GAP are in bold.

Use should be crossed out when the applicant no longer supports this use.

GAP rev. 1, date: 15/07/2025

PPP (product name/code): BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Formulation type: WP (a, b)

Active substance 1: copper (Bordeaux mixture)

Conc. of a.s. 1: 200 g/kg (c)

Applicant: INDUSTRIAS QUIMICAS DEL VALLES S.A.

Professional use: ☒

Zone(s): Southern Zone & Interzonal (d)

Non-professional use: ☐

Verified by MS: Yes

Field of use: Fungicide and bactericide

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergis per ha (^(f))
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
Zonal uses (field or outdoor uses, certain types of protected crops)													
4	FR (SEZ)	Lemon <i>(Citrus limon;</i> 0110030; FC 0002, FC 0204)	F	Phytophthora <i>(Phytophthora spp.;</i> 1PHYTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	Not acceptable (worker,PPE*)
8	FR (SEZ)	Lemon <i>(Citrus limon;</i> 0110030; FC 0002, FC 0204)	F	Blast or black pit <i>(Pseudomonas syringae</i> <i>pv. syringae;</i> PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
10	FR (SEZ)	Lemon (<i>Citrus limon</i> ; 0110030; FC 0002, FC 0204)	F	Brown spot (<i>Alternaria alternata</i> f. sp. <i>citri</i> ; ALTEAC)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
14	FR (SEZ)	Mandarin (<i>Citrus reticulata</i> ; 0110050; FC 0003, FC 0206)	F	Phytophthora (<i>Phytophthora</i> spp.; 1PHYTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	Not acceptable (worker,PPE*)
18	FR (SEZ)	Mandarin (<i>Citrus reticulata</i> ; 0110050; FC 0003, FC 0206)	F	Blast or black pit (<i>Pseudomonas syringae</i> pv. <i>syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
20	FR (SEZ)	Mandarin (<i>Citrus reticulata</i> ; 0110050; FC 0003, FC 0206)	F	Brown spot (<i>Alternaria alternata</i> f. sp. <i>citri</i> ; ALTEAC)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
24	FR (SEZ)	Orange (<i>Citrus</i> sp; 0110020; FC 0004, FC 0208)	F	Phytophthora (<i>Phytophthora</i> spp.; 1PHYTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
28	FR (SEZ)	Orange (<i>Citrus</i> sp; 0110020; FC 0004, FC 0208)	F	Blast or black pit (<i>Pseudomonas syringae</i> pv. <i>syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	Not acceptable (worker,PPE*)
30	FR (SEZ)	Orange (<i>Citrus</i> sp; 0110020; FC 0004, FC 0208)	F	Brown spot (<i>Alternaria alternata</i> f. sp. <i>citri</i> ; ALTEAC)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
34	FR (SEZ)	Grapefruit (<i>Citrus paradise</i> ; 0110010; FC 0005, FC 0203)	F	Phytophthora (<i>Phytophthora</i> spp.; 1PHYTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	Not acceptable (worker,PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
38	FR (SEZ)	Grapefruit (<i>Citrus paradise</i> ; 0110010; FC 0005, FC 0203)	F	Blast or black pit (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
40	FR (SEZ)	Grapefruit (<i>Citrus paradise</i> ; 0110010; FC 0005, FC 0203)	F	Brown spot (<i>Alternaria alternata</i> f. sp. <i>citri</i> ; ALTEAC)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
43	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Leaf curl (<i>Taphrina deformans</i> ; TAPHDE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	Not acceptable (worker,PPE*)
46	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Monilia (<i>Monilia</i> spp.; 1MONLG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
49	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Shot-hole (<i>Stigmina carpophila</i> ; STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
52	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
54	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Cytospora canker (<i>Cytospora leucostoma</i> ; VALSLE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
60	FR (SEZ)	Cherry (<i>Prunus avium</i> ; 0140020; FS 0013, FS0243, FS 0244)	F	Monilia (<i>Monilia</i> spp.; 1MONLG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	Not acceptable (worker,PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
63	FR (SEZ)	Cherry <i>(Prunus avium;</i> 0140020; FS 0013, FS0243, FS 0244)	F	Shot-hole <i>(Stigmina carpophila;</i> STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
66	FR (SEZ)	Cherry <i>(Prunus avium;</i> 0140020; FS 0013, FS0243, FS 0244)	F	Bacterial canker <i>(Pseudomonas syringae</i> <i>pv. syringae;</i> PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
70	FR (SEZ)	Cherry <i>(Prunus avium;</i> 0140020; FS 0013, FS0243, FS 0244)	F	Anthraxnose <i>(Blumeriella jaapii;</i> BLUMJA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
73	FR (SEZ)	Peach <i>(Prunus persica;</i> 0140030; FS 2001, FS 0247)	F	Leaf curl <i>(Taphrina deformans;</i> TAPHDE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	Not acceptable (worker,PPE*)
		Nectarine <i>(Prunus persica var.</i> <i>nucipersica;</i> 0140030; FS 2001, FS 0245)											
76	FR (SEZ)	Peach <i>(Prunus persica;</i> 0140030; FS 2001, FS 0247)	F	Monilia <i>(Monilia spp.;</i> 1MONLG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
		Nectarine <i>(Prunus persica var.</i> <i>nucipersica;</i> 0140030; FS 2001, FS 0245)											

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
79	FR (SEZ)	Peach <i>(Prunus persica;</i> 0140030; FS 2001, FS 0247) Nectarine <i>(Prunus persica var.</i> <i>nucipersica;</i> 0140030; FS 2001, FS 0245)	F	Shot-hole <i>(Stigmia carpophila;</i> STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
82	FR (SEZ)	Peach <i>(Prunus persica;</i> 0140030; FS 2001, FS 0247) Nectarine <i>(Prunus persica var.</i> <i>nucipersica;</i> 0140030; FS 2001, FS 0245)	F	Bacterial canker <i>(Pseudomonas syringae</i> <i>pv. syringae;</i> PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
84	FR (SEZ)	Peach <i>(Prunus persica;</i> 0140030; FS 2001, FS 0247) Nectarine <i>(Prunus persica var.</i> <i>nucipersica;</i> 0140030; FS 2001, FS 0245)	F	Cytospora canker <i>(Cytospora leucostoma;</i> VALSLE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
87	FR (SEZ)	Plum <i>(Prunus sp;</i> 0140040; FS 0014)	F	Leaf curl <i>(Taphrina deformans;</i> TAPHDE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	Not acceptable (worker,PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
90	FR (SEZ)	Plum (<i>Prunus sp</i> ; 0140040; FS 0014)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
93	FR (SEZ)	Plum (<i>Prunus sp</i> ; 0140040; FS 0014)	F	Shot-hole (<i>Stigmia carpophila</i> ; STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
96	FR (SEZ)	Plum (<i>Prunus sp</i> ; 0140040; FS 0014)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
101	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	Scab (<i>Venturia inaequalis</i> ; VENTIN)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker,PPE*)
104	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	European canker (<i>Nectria galligena</i> ; NECTGA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
107	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
110	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	Blister spot (<i>Pseudomonas syringae</i> <i>pv. papulans</i> ; PSDMPA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
112	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	Fire blight (<i>Erwinia amylovora</i> ; ERWIAM)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
115	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	Scab (<i>Venturia pyrina</i> ; VENTPI)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker,PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
118	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	European canker (<i>Nectria galligena</i> ; NECTGA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
121	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
124	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
126	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	Fire blight (<i>Erwinia amylovora</i> ; ERWIAM)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
129	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	Scab (<i>Venturia pyrina</i> ; VENTPI)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker,PPE*)
132	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	European canker (<i>Nectria galligena</i> ; NECTGA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
135	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
138	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
140	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	Fire blight (<i>Erwinia amylovora</i> ; ERWIAM)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
143	FR (SEZ)	Medlar and Japa- nese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	Scab (<i>Venturia inaequalis</i> ; VENTIN)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker,PPE*)
146	FR (SEZ)	Medlar and Japa- nese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	European canker (<i>Nectria galligena</i> ; NECTGA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
149	FR (SEZ)	Medlar and Japa- nese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
152	FR (SEZ)	Medlar and Japa- nese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	Bacterial canker (<i>Pseudomonas syringae pv. syringae</i> ; PSDMSY)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
154	FR (SEZ)	Medlar and Japa- nese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	Fire blight (<i>Erwinia amylovora</i> ; ERWIAM)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
157	FR (SEZ)	Almond (<i>Prunus dulcis</i> ; 0120010; TN 0660)	F	Leaf curl (<i>Taphrina deformans</i> ; TAPHDE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker,PPE*)
160	FR (SEZ)	Almond (<i>Prunus dulcis</i> ; 0120010; TN 0660)	F	Monilia (<i>Monilia spp.</i> ; 1MONIG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
163	FR (SEZ)	Almond (<i>Prunus dulcis</i> ; 0120010; TN 0660)	F	Shot-hole (<i>Stigmina carpophila</i> ; STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
166	FR (SEZ)	Almond (<i>Prunus dulcis</i> ; 0120010; TN 0660)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
172	FR (SEZ)	Walnut (<i>Juglans regia</i> ; 0120110; TN 0678)	F	Alternaria (<i>Alternaria spp.</i> ; 1ALTEG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker,PPE*)
177	FR (SEZ)	Walnut (<i>Juglans regia</i> ; 0120110; TN 0678)	F	Bacterial blight (<i>Xanthomonas arboricola</i> <i>pv. juglandis</i> ; XANTJU)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
180	FR (SEZ)	Walnut (<i>Juglans regia</i> ; 0120110; TN 0678)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
184	FR (SEZ)	Hazelnut (<i>Corylus avellana</i> ; 0120060; TN 0666)	F	Alternaria (<i>Alternaria</i> spp.; 1ALTEG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker,PPE*)
186	FR (SEZ)	Hazelnut (<i>Corylus avellana</i> ; 0120060; TN 0666)	F	Cytospora canker (<i>Cytospora</i> spp; 1CYTOG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
189	FR (SEZ)	Hazelnut (<i>Corylus avellana</i> ; 0120060; TN 0666)	F	Bacterial blight (<i>Xanthomonas arboricola</i> pv. <i>corylina</i> ; XANTCY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
192	FR (SEZ)	Hazelnut (<i>Corylus avellana</i> ; 0120060; TN 0666)	F	Stem dieback (<i>Pseudomonas avellanae</i> ; PSDMAL)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
194	FR (SEZ)	Chestnut (<i>Castanea sativa</i> ; 0120040; TN 0664)	F	Mycosphaerella (<i>Mycosphaerella</i> spp.; 1MYCOG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker,PPE*)
196	FR (SEZ)	Chestnut (<i>Castanea sativa</i> ; 0120040; TN 0664)	F	Alternaria (<i>Alternaria</i> spp.; 1ALTEG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
201	FR (SEZ)	Chestnut (<i>Castanea sativa</i> ; 0120040; TN 0664)	F	Xanthomonas (<i>Xanthomonas</i> spp; 1XANTG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
204	FR (SEZ)	Chestnut (<i>Castanea sativa</i> ; 0120040; TN 0664)	F	Pseudomonas (<i>Pseudomonas</i> spp; 1PSDMG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
208	FR (SEZ)	Pistachio (<i>Pistacia vera</i> ; 0120100; TN 0675)	F	Alternaria (<i>Alternaria</i> spp.; 1ALTEG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker,PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
213	FR (SEZ)	Pistachio (<i>Pistacia vera</i> ; 0120100; TN 0675)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
216	FR (SEZ)	Pistachio (<i>Pistacia vera</i> ; 0120100; TN 0675)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
220	FR (SEZ)	Olive (Table olives, olive for oil production) (<i>Olea europaea</i> ; 0161030, 0402010; FT 0305, SO 0305)	F	Peacock spot (<i>Spilocaea oleaginea</i> ; CYCLOL)	Foliar spray	10-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	600- 1000	14	Not acceptable (worker,PPE*)
224	FR (SEZ)	Olive (Table olives, olive for oil production) (<i>Olea europaea</i> ; 0161030, 0402010; FT 0305, SO 0305)	F	Bacterial canker (<i>Pseudomonas savastoni</i> <i>pv. savastoni</i> ; PSDMSA)	Foliar spray	10-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	600- 1000	14	
228	FR (SEZ)	Olive (Table olives, olive for oil production) (<i>Olea europaea</i> ; 0161030, 0402010; FT 0305, SO 0305)	F	Anthracnose (<i>Colletotrichum</i> <i>gloeosporioides</i> ; GLOMCI)	Foliar spray	10-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	600- 1000	14	
230	FR (SEZ)	Grape (Table grape, wine grapes) (<i>Vitis vinifera</i> ; 0151000, 0151010, 0151020; FB 0269)	F	Downy mildew (<i>Plasmopara viticola</i> ; PLASVI)	Foliar spray	15-81 & 91	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	21	Not acceptable (worker,PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
234	FR (SEZ)	Grape (Table grape, wine grapes) <i>(Vitis vinifera</i> ; 0151000, 0151010, 0151020; FB 0269)	F	Bacterial blight <i>(Xanthomonas ampelina</i> ; XANTAM)	Foliar spray	15-81 & 91	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	21	
237	FR (SEZ)	Tomato <i>(Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Late blight <i>(Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	Not acceptable (PPE*)
240	FR (SEZ)	Tomato <i>(Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Early blight <i>(Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
243	FR (SEZ)	Tomato <i>(Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Anthraxnose <i>(Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
246	FR (SEZ)	Tomato <i>(Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Bacterial spot <i>(Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
249	FR (SEZ)	Tomato <i>(Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Bacterial speck <i>(Pseudomonas syringae pv. tomato</i> ; PSDMTM)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
252	FR (SEZ)	Aubergine (Eggplant) <i>(Solanum melongena</i> ; 0231030; VO 0440)	F	Late blight <i>(Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	Not acceptable (PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
255	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	F	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
258	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	F	Anthrachnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
261	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	F	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
264	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	F	Bacterial speck (<i>Pseudomonas syringae pv. tomato</i> ; PSDMTM)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
267	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Blight of pepper (<i>Phytophthora capsici</i> ; PHYTCP)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRI,PPE*)
270	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
273	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Anthrachnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (^(f))
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
276	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
279	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
280	FR (SEZ)	Leek (<i>Allium porrium</i> ; 0270060; VA 0384) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Downy mildew (<i>Phytophthora porri</i> ; PHYTPO)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL ,PPE*)
287	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Downy mildew (<i>Bremia lactucae</i> ; BREMLA)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL ,PPE*)
290	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Ascochyta (<i>Ascochyta spp</i> ; (1ASCOG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
293	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
296	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
299	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
300	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Downy mildew (<i>Plasmopara crustosa</i> ; PLASCR)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL ,PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
301	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Ascochyta (<i>Ascochyta spp</i> ; (1ASCOG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
302	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
303	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
304	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
305	FR (SEZ)	Cardoon (<i>Cynara carduncu- lus</i> ; 0270020; VS 0623) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Downy mildew (<i>Bremia lactucae</i> ; BREMLA)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL ,PPE*)
306	FR (SEZ)	Cardoon (<i>Cynara carduncu- lus</i> ; 0270020; VS 0623) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Ascochyta (<i>Ascochyta spp</i> ; 1ASCOG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
307	FR (SEZ)	Cardoon (<i>Cynara carduncu- lus</i> ; 0270020; VS 0623) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
308	FR (SEZ)	Cardoon (<i>Cynara carduncu- lus</i> ; 0270020; VS 0623) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
309	FR (SEZ)	Cardoon (<i>Cynara carduncu- lus</i> ; 0270020; VS 0623) Whole group of stem vegetables ex- cept asparagus (0270000; VS 0078)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
312	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	Downy mildew (<i>Hyaloperonospora brassicae</i> ; HPERBR)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL ,PPE*)
315	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	Black spot (<i>Alternaria brassicae</i> ; ALTEBA)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
318	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	<i>Colletotrichum</i> (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
321	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	Black rot (<i>Xanthomonas campestris pv. campestris</i> ; XANTCA)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
324	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
327	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Downy mildew (<i>Hyaloperonospora</i> <i>brassicae</i> ; HPERBR)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL ,PPE*)
330	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Black spot (<i>Alternaria brassicae</i> ; ALTEBA)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
333	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Colletotrichum (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
336	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Black rot (<i>Xanthomonas campestris</i> pv. <i>campestris</i> ; XANTCA)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
339	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
342	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not Acceptable (PPE*)
345	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
348	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
351	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
354	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
357	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not Acceptable (PPE*)
360	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
363	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Anthraxnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
366	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
369	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
372	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not Acceptable (PPE*)
375	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
378	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
381	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
384	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
387	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*)
390	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (open- field and walk-in tunnel)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
393	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
396	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
399	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
402	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*)
405	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
408	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
411	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
414	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
417	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
420	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
423	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Anthraco nose (<i>Colletotrichum lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
426	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Bacterial leaf spot (<i>Xanthomonas campestris pv. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
429	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Angular leaf spot (<i>Pseudomonas syringae pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
432	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	Downy mildew (<i>Peronospora destructor</i> ; PERODE)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
435	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	Purple blotch (<i>Alternaria porri</i> ; ALTEPO)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
438	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	Anthraco nose (<i>Colletotrichum coccodes</i> ; COLLCC)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
440	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	Leaf blight (<i>Pleospora allii</i> ; PLEOAL)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
443	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
446	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	<i>Xanthomonas</i> (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
449	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	Downy mildew (<i>Peronospora destructor</i> ; PERODE)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
452	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	Purple blotch (<i>Alternaria porri</i> ; ALTEPO)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
455	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	Antracnose (<i>Colletotrichum coccodes</i> ; COLLCC)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
457	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	Leaf blight (<i>Pleospora allii</i> ; PLEOAL)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
460	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
463	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	<i>Xanthomonas</i> (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
466	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Downy mildew (<i>Peronospora destructor</i> ; PERODE)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
469	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Purple blotch (<i>Alternaria porri</i> ; ALTEPO)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
472	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Antracnose (<i>Colletotrichum coccodes</i> ; COLLCC)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
474	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Leaf blight (<i>Pleospora allii</i> ; PLEOAL)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
477	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
480	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
483	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	F	Downy mildew (<i>Bremia lactucae</i> ; BREMLA)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*) Not acceptable against grey mould and sclerotinia (efficacy)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
486	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	F	Leaf spot (<i>Alternaria sonchi</i> ; ALTESN)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
489	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	F	Anthracnose (<i>Marssonina panattoniana</i> ; MARSPA)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
491	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	F	Sclerotinia (<i>Sclerotinia minor</i> ; SCLEMI)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
494	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
497	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	F	Bacterial spot (<i>Xanthomonas axonopodis</i> pv. <i>vitians</i> ; XANTVI)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
499	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and simi- lar (leaves) (0252000)	F	Downy mildew (<i>Peronospora farinosa</i> f. sp. <i>Spinaciae</i> ;PEROFS)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*)
501	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and simi- lar (leaves) (0252000)	F	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
503	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and simi- lar (leaves) (0252000)	F	Anthrachnose (<i>Colletotrichum spinaciae</i> ; COLLDS)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
510	FR (SEZ)	Bean (Bean with pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Downy mildew (<i>Phytophthora phaseoli</i> ; PHYTPH)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not Acceptable against rust (efficacy)
513	FR (SEZ)	Bean (Bean with pods, bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Anthrachnose (<i>Colletotrichum lindemuthianum</i> ; COLLLD)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
515	FR (SEZ)	Bean (Bean with pods, bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Rust (<i>Uromyces appendiculatus</i> ; UROMAP)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
517	FR (SEZ)	Bean (Bean with pods, bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Septoria (<i>Septoria spp</i> ; 1SEPTG)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
520	FR (SEZ)	Bean (Bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Bacterial blight (<i>Xanthomonas axonopodis</i> pv. <i>phaseoli</i> ; XANTPH)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
523	FR (SEZ)	Bean (Bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
526	FR (SEZ)	Pea (Pea with pods) (<i>Pisum sativum</i> ; 0260030, 0260040; VP 4441)	F	Downy mildew (<i>Peronospora viciae</i> ; PEROVI)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (MRL,PPE*) Not acceptable against rust (efficacy)
529	FR (SEZ)	Pea (Pea with pods, pea without pods) (<i>Pisum sativum</i> ; 0260030, 0260040; VP 4441)	F	Anthraco nose (<i>Colletotrichum</i> <i>lindemuthianum</i> ; COLLLD)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
531	FR (SEZ)	Pea (Pea with pods, pea without pods) (<i>Pisum sativum</i> ; 0260030, 0260040; VP 4441)	F	Rust (<i>Uromyces fabae</i> ; UROMVF)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
533	FR (SEZ)	Pea (Pea with pods, pea without pods) (<i>Pisum sativum</i> ; 0260030, 0260040; VP 4441)	F	Leaf blotch (<i>Septoria pisi</i> ; SEPTPI)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
545	FR (SEZ)	Lentils (<i>Lens culinaris</i> ; 0260050; VP 0533)	F	Anthrachnose (<i>Colletotrichum lindemuthianum</i> ; COLLLD)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
547	FR (SEZ)	Lentils (<i>Lens culinaris</i> ; 0260050; VP 0533)	F	Rust (<i>Uromyces fabae</i> ; UROMVF)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (MRL,PPE*) Not acceptable against rust (efficacy)
549	FR (SEZ)	Lentils (<i>Lens culinaris</i> ; 0260050; VP 0533)	F	Septoria (<i>Septoria spp</i> ; 1SEPTG)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
558	FR (SEZ)	Potato (<i>Solanum tuberosum</i> ; 0211000; VR 0589)	F	Late blight (<i>Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL,PPE*)
561	FR (SEZ)	Potato (<i>Solanum tuberosum</i> ; 0211000; VR 0589)	F	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
563	FR (SEZ)	Potato (<i>Solanum tuberosum</i> ; 0211000; VR 0589)	F	Anthrachnose (<i>Colletotrichum coccodes</i> ; COLLCC)	Foliar spray	15-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
570	FR (SEZ)	Carrot (<i>Daucus carota subsp. Sativus</i> ; 0213020; VR 0577) Other root and tu- ber vegetables except sugar beet (0213000)	F	Downy mildew (<i>Plasmopara nivea</i> ; PLASCR)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL,PPE*)
571	FR (SEZ)	Carrot (<i>Daucus carota subsp. Sativus</i> ; 0213020; VR 0577) Other root and tu- ber vegetables except sugar beet (0213000)	F	Leaf blight (<i>Alternaria dauci</i> ; ALTEDA)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
572	FR (SEZ)	Carrot (<i>Daucus carota subsp. Sativus</i> ; 0213020; VR 0577) Other root and tu- ber vegetables except sugar beet (0213000)	F	Anthrachnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
573	FR (SEZ)	Carrot (<i>Daucus carota</i> <i>subsp. Sativus</i> ; 0213020; VR 0577) Other root and tu- ber vegetables except sugar beet (0213000)	F	Brown leaf spot (<i>Cercospora carotae</i> ; CERCCA)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
574	FR (SEZ)	Carrot (<i>Daucus carota</i> <i>subsp. Sativus</i> ; 0213020; VR 0577) Other root and tu- ber vegetables except sugar beet (0213000)	F	Bacterial leaf blight (<i>Xanthomonas campestris</i> <i>pv. carotae</i> ; XANTCR)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
575	FR (SEZ)	Carrot (<i>Daucus carota</i> <i>subsp. Sativus</i> ; 0213020; VR 0577) Other root and tu- ber vegetables except sugar beet (0213000)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
576	FR (SEZ)	Sugar beet (<i>Beta vulgaris subsp.</i> <i>vulgaris</i> ; 0900010; VR 0596)	F	Leaf spot (<i>Cercospora beticola</i> ; CERCBE)	Foliar spray	39-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL,PPE*)
579	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Anthracnose (<i>Colletotrichum fragariae</i> ; COLLFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
580	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Leaf spot (<i>Mycosphaerella fragariae</i> ; MYCOFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
581	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Red stele (<i>Phytophthora fragariae</i> ; PHYTFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
582	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Angular leaf spot (<i>Xanthomonas fragariae</i> ; XANTFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
583	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
585	FR (SEZ)	Ornamental plants and trees	F	Peronospora (<i>Peronospora spp</i> ; 1PEROG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	Not acceptable (worker,PPE*) Not acceptable against rust (efficacy)
588	FR (SEZ)	Ornamental plants and trees	F	Septoria (<i>Septoria spp</i> ; 1SEPTG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
590	FR (SEZ)	Ornamental plants and trees	F	Colletotrichum (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
593	FR (SEZ)	Ornamental plants and trees	F	Puccinia (<i>Puccinia spp</i> ; 1PUCCG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
595	FR (SEZ)	Ornamental plants and trees	F	Monilia (<i>Monilia spp</i> ; 1MONLG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
Interzonal uses (use as seed treatment, in greenhouses (or other closed places of plant production), as post-harvest treatment or for treatment of empty storage rooms)													
598	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Late blight (<i>Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
601	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
604	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Anthraxnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
607	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
610	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Bacterial speck (<i>Pseudomonas syringae pv. tomato</i> ; PSDMTM)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
613	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Late blight (<i>Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
616	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
619	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Anthraxnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
622	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
625	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Bacterial speck (<i>Pseudomonas syringae pv. tomato</i> ; PSDMTM)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
627	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	G	Blight of pepper (<i>Phytophthora capsici</i> ; PHYTCP)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL,PPE*)
629	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	G	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
631	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	G	Anthracnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
633	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	G	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
635	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	G	Bacterial canker (<i>Pseudomonas syringae pv. syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
638	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Downy mildew (<i>Pseudoperonospora cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
641	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
644	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Anthraxnose (<i>Colletotrichum lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
647	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Bacterial leaf spot (<i>Xanthomonas campestris pv. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
650	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Angular leaf spot (<i>Pseudomonas syringae pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
653	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Downy mildew (<i>Pseudoperonospora cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
656	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
659	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
662	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
665	FR (SEZ)	Courgette (zuc- chini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
668	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
671	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
674	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
677	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
680	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
682	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*)
684	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
686	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Anthraxnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
688	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
690	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
692	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*)
694	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
696	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
698	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
700	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
702	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*)
704	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
706	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
708	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
710	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (^(f))
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
713	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	G	Downy mildew (<i>Bremia lactucae</i> ; BREMLA)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (PPE*) Not acceptable against grey mould and sclerotinia (efficacy)
716	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	G	Leaf spot (<i>Alternaria sonchi</i> ; ALTESN)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
719	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	G	Anthrachnose (<i>Marssonina panattoniana</i> ; MARSPA)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
721	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	G	Sclerotinia (<i>Sclerotinia minor</i> ; SCLEMI)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
724	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	G	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
727	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants includ- ing leaves and sprouts of brassica- cea (0251000)	G	Bacterial spot (<i>Xanthomonas axonopodis</i> <i>pv. vitians</i> ; XANTVI)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
729	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and simi- lar (leaves) (0252000)	G	Downy mildew (<i>Peronospora farinosa</i> f. <i>sp. Spinaciae</i> ;PEROFS)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*)
731	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and simi- lar (leaves) (0252000)	G	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
733	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and simi- lar (leaves) (0252000)	G	Anthrachnose (<i>Colletotrichum spinaciae</i> ; COLLDS)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
739	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	Anthrachnose (<i>Colletotrichum fragariae</i> ; COLLFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL,PPE*))
740	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	Leaf spot (<i>Mycosphaerella</i> <i>fragariae</i> ; MYCOFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
741	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	Red stele (<i>Phytophthora fragariae</i> ; PHYTFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
742	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	Angular leaf spot (<i>Xanthomonas fragariae</i> ; XANTFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
743	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
745	FR (SEZ)	Ornamental plants and trees	G	<i>Peronospora</i> (<i>Peronospora spp</i> ; 1PEROG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	Not acceptable (worker,PPE*) Not acceptable against rust (efficacy)
748	FR (SEZ)	Ornamental plants and trees	G	<i>Septoria</i> (<i>Septoria spp</i> ; 1SEPTG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
750	FR (SEZ)	Ornamental plants and trees	G	<i>Colletotrichum</i> (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
753	FR (SEZ)	Ornamental plants and trees	G	<i>Puccinia</i> (<i>Puccinia spp</i> ; 1PUCCG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
755	FR (SEZ)	Ornamental plants and trees	G	<i>Monilia</i> (<i>Monilia spp</i> ; 1MONLG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	

* PPE : Personal protective equipment. The information is insufficient to ensure the performance of the clothing PPE (in accordance with standard NF EN ISO 27065/A1) for this type of formulation (WP). A. Shaw *and al.* (2018).

**Remarks
table
heading:**

- (a) e.g. wettable powder (WP), emulsifiable concentrate (EC), granule (GR)
 (b) Catalogue of pesticide formulation types and international coding system CropLife
 International Technical Monograph n°2, 6th Edition Revised May 2008
 (c) g/kg or g/l

- (d) Select relevant
 (e) Use number(s) in accordance with the list of all intended GAPs in Part B, Section 0 should be given
 in column 1
 (f) No authorisation possible for uses where the line is highlighted in grey, Use should be crossed out
 when the notifier no longer supports this use.

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K

Part A - National Assessment

FRANCE

Remarks	1	Numeration necessary to allow references	7	Growth stage at first and last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application
	2	Use official codes/nomenclatures of EU Member States	8	The maximum number of application possible under practical conditions of use must be provided.
columns:	3	For crops, the EU and Codex classifications (both) should be used; when relevant, the use situation should be described (e.g. fumigation of a structure)	9	Minimum interval (in days) between applications of the same product
	4	F: professional field use, Fn: non-professional field use, Fpn: professional and non-professional field use, G: professional greenhouse use, Gn: non-professional greenhouse use, Gpn: professional and non-professional greenhouse use, I: indoor application	10	For specific uses other specifications might be possible, e.g.: g/m ³ in case of fumigation of empty rooms. See also EPPO-Guideline PP 1/239 Dose expression for plant protection products.
	5	Scientific names and EPPO-Codes of target pests/diseases/ weeds or, when relevant, the common names of the pest groups (e.g. biting and sucking insects, soil born insects, foliar fungi, weeds) and the developmental stages of the pests and pest groups at the moment of application must be named.	11	The dimension (g, kg) must be clearly specified. (Maximum) dose of a.s. per treatment (usually g, kg or L product/ha).
	6	Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plants - type of equipment used must be indicated.	12	If water volume range depends on application equipments (e.g. ULVA or LVA) it should be mentioned under "application: method/kind".
			13	PHI - minimum pre-harvest interval
			14	Remarks may include: Extent of use/economic importance/restrictions

3 Background of authorisation decision and risk management

3.1 Physical and chemical properties (Part B, Section 2)

BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) is a wettable powder formulation (WP and WP-SB). All studies have been performed in accordance with the current requirements and the results are deemed acceptable. The appearance of the product is an odourless fine green powder formulation. The product is not explosive and has no oxidising properties. It is not flammable. It has not a self-ignition temperature below 400°C. In aqueous solution 1%, it has a pH value of 6.75 at 22°C. There is no effect of high temperature on the stability of the formulation, since after 14 days at 54 °C, neither the active ingredient content nor the technical properties were changed. The stability data indicate a shelf life of at least two years at ambient temperature when stored in LDPE and HDPE, PE or PET (water soluble bags). Its technical characteristics are acceptable for a WP formulation.

The formulation is not classified for the physico-chemical aspect.

When water soluble bags are used, the slurry should be stirred at least for 3 minutes before application.

3.2 Efficacy (Part B, Section 3)

Considering the data submitted:

- The efficacy level of BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) is considered acceptable for most of the requested uses. For uses intended to control grey mould and sclerotinia on lettuce, rusts on beans, peas, floral crops and green plants, and cercosporiosis on beetroot, the demonstration of the efficacy of the product is considered insufficient.
- The phytotoxicity level of BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) is considered as acceptable for all the requested uses. On ornamental crops, considering the diversity of crops, it is recommended, in case of doubts, to test the product selectivity on a small area. Moreover, it is also recommended to avoid applications during flowering, because of a risk of burnings on flowers.
- The risks of negative impact on yield, cider making process, propagation, succeeding crops, adjacent crops are considered as negligible. Risks with copper such as spotting of table grape berries or on the wine-making process are known. However, these risks of negative impact are considered acceptable.
- There is a risk of resistance developing or appearing to copper for *Xanthomonas* bacteria that requires a monitoring on vegetable crops (e.g. tomato) and walnut.

3.3 Methods of analysis (Part B, Section 5)

3.3.1 Analytical method for the formulation

Analytical methods for the determination of the active substance and the relevant impurities in the formulation are available and validated.

3.3.2 Analytical methods for residues

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Analytical methods are available in the Draft Assessment Report (DAR) and in this dossier, and are validated for the determination of residues of copper in plants (high water, oily, acidic and dry content commodities), soil, water (surface and drinking), air and body fluids.

According to EFSA conclusions, an ILV of the analytical methods for the determination of residues of copper in plants is required.

Analytical methods for the determination of residues of copper in foodstuffs of animal origin are missing and are required. Moreover, the LOQ of the available methods for the determination of residues of copper in water is not in accordance with the European Directive 98/83/EC.

3.4 Mammalian toxicology (Part B, Section 6)

Endpoints used in risk assessment

Agreed EU endpoints	
Active substance	Copper (Bordeaux mixture) 200 g/kg
AOEL systemic	0.08 mg/kg bw/d
AAOEL	Not necessary
Inhalation absorption	100%
Oral absorption	50%
Vapour pressure	Not necessary
Dermal absorption	Concentrate: 1% Dilution: 9% (Based on <i>in vitro</i> through human skin studies)

3.4.1 Acute toxicity

BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC), containing 200 g/kg copper (Bordeaux mixture), has a low acute oral and dermal toxicity, is harmful if inhaled, and is not irritating to the rabbit skin or eye and is not a skin sensitiser.

3.4.2 Operator exposure

Considering proposed uses, operator systemic exposure was estimated using EFSA model¹²:

Outdoor:

		Copper under the form of Bordeaux mixture	
Model data	Level of PPE	Total absorbed dose (mg/kg/day)	% of systemic AOEL
Critical use: Olive (dense foliage)			
Tractor mounted outdoor, upward applications			

¹² AOEM – Agricultural Operator Exposure Model (EFSA Journal 2014;12 (10):3874)

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.0976	121.99
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0437	54.62
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0309	38.67
Manual hand held outdoor, upward applications			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.1095	136.84
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0823	102.91
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP2, P2 and similar M/L	0.0769	96.12
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0766	95.73
Critical use: Grape			
Tractor mounted outdoor, Upward applications			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.0976	121.99
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0437	54.62
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0309	38.67
Manual knapsack outdoor, Upward applications			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves M/L and A	0.0036	4.48
Critical use: Strawberry (Low berries and other small fruits).			

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Tractor mounted outdoor, Downward applications			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.1257	157.15
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.3856	48.21
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0136	16.97
Manual hand held outdoor, Downward applications			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.0923	115.71
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0515	64.44
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0424	53.01
Critical use: Fruiting vegetable (covers Brassica vegetable, Bulb vegetable, legume, root and tuber vegetable)			
Tractor mounted outdoor, Downward applications			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.1257	157.15
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0386	48.21
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.01358	16.97
Manual hand held outdoor, Downward applications			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.092	115.71
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0515	64.44
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0424	53.01

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

	covered) and gloves M/L and A + water soluble bags M/L		
Critical use: Stone fruits (covers Pome fruits, Tree nuts and Citrus). Dense foliage			
Tractor mounted outdoor, upward applications			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.0976	121.99
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0437	54.62
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0309	38.67
Manual hand held outdoor, upward applications			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.1095	136.84
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0823	102.91
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP2, P2 and similar M/L	0.0769	96.12
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0766	95.73
Critical use: Ornamentals			
Tractor mounted outdoor, upward applications			
Application rate		1 kg a.s./ha, 3 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.0976	121.99
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0437	54.62
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0309	38.67
Manual hand held outdoor, upward applications			
Application rate		1 kg a.s./ha, 3 applications	
Spray application EFSA model	Work wear (arms, body and legs covered) and gloves M/L and Application	0.0620	77.54

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Body weight: 60 kg	Application		
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0119	14.84

Indoor:

Copper under the form of Bordeaux mixture			
Model data	Level of PPE	Total absorbed dose (mg/kg/day)	% of systemic AOEL
Critical use: Strawberry (Low berries and other small fruits).			
Manual hand held, upward spraying			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.0620	77.54
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0210	26.27
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0119	14.84
Fruiting vegetable: tomato (critical use)			
Manual knapsack , upward spraying			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves M/L and A	0.0036	4.48
Manual hand held, upward spraying			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.0620	77.54
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0119	14.84
Critical use: Leaf vegetable			
Manual hand held , downward spraying			
Application rate		1 kg a.s./ha, 4 applications	
Spray application EFSA model	Work wear (arms, body and legs covered) and gloves	0.0923	115.71

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Body weight: 60 kg	M/L and Application		
	Work wear (arms, body and legs covered) and gloves M/L/A, mask (FP1, P1 and similar) during M/L	0.051	64.44
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0424	53.01
Critical use: Ornamentals			
Manual knapsack, upward spraying			
Manual hand held, upward spraying.			
Application rate		1 kg a.s./ha, 3 applications	
Spray application EFSA model Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and Application	0.062	77.54
	Work wear (arms, body and legs covered) and gloves M/L and Application + FP1, P1 and similar M/L	0.0210	26.27
	Work wear (arms, body and legs covered) and gloves M/L and A + water soluble bags M/L	0.0119	14.84

Conclusion:

According to the EFSA model calculations, it can be concluded that the risk for the operator using BM 20 WP NC is **acceptable** taking into account the above mentioned personal protective equipment (PPE). The packaging “water soluble bag” permits to avoid the recommendation of RPE.

For details of personal protective equipment for operators, refer to the Decision in Appendix 1.

3.4.3 Worker exposure

Workers may have to enter treated areas after treatment for crop inspection/irrigation or cutting, sorting, bundling, carrying or hand harvesting or maintenance or reaching, picking or searching, reaching, picking activities. Therefore, estimation of worker exposure was calculated according to AOEM model.

Outdoor:

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
 Part A - National Assessment
 FRANCE

Model data		Bordeaux mixture
	Level of PPE	%AOEL
Oilfruits Activity: Searching , reaching, picking Outdoor Work rate: 8 hours/day Number of applications : 4 Interval between treatments: 7 days		
DT50:		30 days
DFR:		3 µg/cm ² /kg a.s./ha
Application rate (kg as/ha)		1 kg Bordeaux mixture/ha
Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves TC: 2250 cm ² /person/h	189.84
Grapes Activity: hand harvesting Outdoor Work rate: 8 hours/day Number of applications : 4 Interval between treatments: 7 days		
DT50:		30 days
DFR:		3 µg/cm ² /kg a.s./ha
Application rate (kg as/ha)		1 kg Bordeaux mixture/ha
Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves TC: xxx cm ² /person/h	852.19
Ornamentals Activity: cutting, sorting, bundling, carrying Outdoor Work rate: 8 hours/day Number of applications : 3 Interval between treatments: 7 days		
DT50:		30 days
DFR:		3 µg/cm ² /kg a.s./ha
Application rate (kg as/ha)		1 kg Bordeaux mixture/ha
Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves TC: 1400 cm ² /person/h	110.25

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Strawberry Activity: Reaching, picking Outdoor Work rate: 8 hours/day Number of applications : 4 Interval between treatments: 7 days		
DT50:		30 days
DFR:		3 µg/cm ² /kg a.s./ha
Application rate (kg as/ha)		1 kg Bordeaux mixture/ha
Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves TC: 750 cm ² /person/h	63.28
Fruiting vegetables Activity: Reaching, picking Outdoor Work rate: 8 hours/day Number of applications : 4 Interval between treatments: 7 days		
DT50:		30 days
DFR:		3 µg/cm ² /kg a.s./ha
Application rate (kg as/ha)		1 kg Bordeaux mixture/ha
Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves TC: 580 cm ² /person/h	48.94
Potato Activity: inspection, irrigation Outdoor Work rate: 8 hours/day Number of applications : 4 Interval between treatments: 7 days		
DT50:		30 days
DFR:		3 µg/cm ² /kg a.s./ha
Application rate (kg as/ha)		1 kg Bordeaux mixture/ha
Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves TC: 1400 cm ² /person/h	29.53

Conclusion for outdoor uses:

It is concluded that there is an unacceptable risk anticipated for the worker for orchards, grapes and ornamentals.

Indoor:

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Model data		Bordeaux mixture
	Level of PPE	%AOEL
Ornamentals Activity: cutting, sorting, bundling, carrying Outdoor Work rate: 8 hours/day Number of applications : 3 Interval between treatments: 7 days		
DT50:		30 days
DFR:		3 µg/cm ² /kg a.s./ha
Application rate (kg as/ha)		1 kg Bordeaux mixture/ha
Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves TC: 1400 cm ² /person/h	126.92
Strawberry Activity: Reaching, picking Outdoor Work rate: 8 hours/day Number of applications : 4 Interval between treatments: 7 days		
DT50:		30 days
DFR:		3 µg/cm ² /kg a.s./ha
Application rate (kg as/ha)		1 kg Bordeaux mixture/ha
Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves TC: 750 cm ² /person/h	63.28
Fruiting vegetables Activity: Reaching, picking Outdoor Work rate: 8 hours/day Number of applications : 4 Interval between treatments: 7 days		
DT50:		30 days
DFR:		3 µg/cm ² /kg a.s./ha
Application rate (kg as/ha)		1 kg Bordeaux mixture/ha
Body weight: 60 kg	Work wear (arms, body and legs covered) + gloves TC: 580 cm ² /person/h	48.94

Conclusion for indoor uses:

There is no unacceptable risk anticipated for worker re-entering treated crop except for ornamentals.

For details of personal protective equipment for workers, refer to the Decision in Appendix 1.

3.4.4 Bystander exposure

Consideration of acute exposure should only be made where an AAOEL has been established during an approval, review or renewal evaluation of an active substance, i.e. no acute operator or bystander exposure assessments can be performed with the AOEM model where no AAOEL has been set¹³.

According to EFSA Guidance on the assessment of exposure of operators, workers, residents and bystanders in risk assessment for plant protection products (EFSA Journal 2014;12(10):3874): “No bystander risk assessment is required for PPPs that do not have significant acute toxicity or the potential to exert toxic effects after a single exposure. Exposure in this case will be determined by average exposure over a longer duration, and higher exposures on one day will tend to be offset by lower exposures on other days. Therefore, exposure assessment for residents also covers bystander exposure.”

No AAOEL has been set for copper. Thus, for this active substance, residents exposure assessment covers bystanders exposure.

3.4.5 Resident exposure

Outdoor: Residential exposure was assessed according to EFSA model incorporating a distance of 10 meters for high crops and 2-3 meters for low crops.

Model data		Bordeaux mixture
		% AOEL
Scenario: Stone fruits (worst case) Buffer zone: 10(m) Drift reduction technology: no Number of applications : 4 Interval between treatments: 7 days		
DT ₅₀		30 days
DFR		3 µg/cm²/kg a.s./h
Resident (children) Body weight: 10 kg	Spray drift (75th percentile)	39.47
	Vapour (75th percentile)	1.34
	Surface deposits (75th percentile)	6.44
	Entry into treated crops (75th percentile)	35.60
	All pathways (mean)	60.09
Resident (adults) Body weight: 60 kg	Spray drift (75th percentile)	21.75
	Vapour (75th percentile)	0.29
	Surface deposits (75th percentile)	2.30
	Entry into treated crops (75th percentile)	19.78
	All pathways (mean)	31.85

An acceptable risk was determined for residents (adult and child) without drift reduction technology.

¹³ Guidance on the assessment of exposure of operators, workers, residents and bystanders in risk assessment for plant protection products (SANTE-10832-2015 rev. 1.7, 2017)

Indoor: in the context of indoor uses, resident exposure is not relevant.

Conclusion:

In conclusion, according to the EFSA model, there is **no unacceptable** risk anticipated for resident (adult and child) after incidental exposure to copper.

3.4.1 Combined exposure

Not relevant. The product contains only one active substance.

3.5 Residues and consumer exposure (Part B, Section 7)

As ornamental plants and trees are non edible commodities, the respective intended uses were not assessed.

For France, an exceedance of the current MRL for copper as laid down in Reg. (EC) 396/2005 of 50 mg/kg in grapes, 5 mg/kg in pome fruits, 5 mg/kg in stone fruits, 30 mg/kg in tree nuts and pistachio, 30 mg/kg in olive, 20 mg/kg in citrus trees, 5.0 mg/kg in outdoor and indoor tomato and aubergine, 5.0 mg/kg in outdoor and indoor edible peel cucurbits, 5.0 mg/kg in onion, garlic and shallot, 5.0 mg/kg in parsnip, Jerusalem artichoke, parsley, celeriac, horseradish and scorzonera, 100 mg/kg in indoor lettuce and other salads and 20 mg/kg in fresh beans and peas with pods and fresh beans without pods is not expected.

Due to MRL exceedance, the uses on strawberries (outdoor and indoor), inedible peel cucurbits (outdoor and indoor), spring onion, spinaches and similar leaves (outdoor and indoor), pepper (outdoor and indoor) and potatoes cannot be recommended.

Due to insufficient residue trials, the uses on flowering brassica, lettuce (outdoor), fresh peas without pods, fresh lentils, carrots, beetroot, leeks, celery, cardoon, and artichokes cannot be recommended.

The acute exposure calculations were not carried out because an acute reference dose (ARfD) was not deemed necessary for copper.

For chronic intake of copper residues, the calculation includes uncertainties linked to the methodology. Therefore, zRMS considers that the risk assessment for consumers cannot be finalised.

zRMS considers no firm conclusion can be reached for any of the intended uses of the product BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC).

Summary for BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC)

Crop	PHI for BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) requested by applicant	PHI/ Withholding period* sufficiently supported for Copper	PHI for BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) Blue proposed by zRMS	zRMS Comments (if different PHI proposed)
Outdoor uses				
Citrus trees	14 days	Yes	14 days	
Stone fruits	F**	Yes	F	
Pome fruits	F	Yes	F	
Tree nuts, pistachio	F	Yes	F	

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
 Part A - National Assessment
 FRANCE

Crop	PHI for BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) requested by applicant	PHI/ Withholding period* sufficiently supported for Copper	PHI for BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) Blue proposed by zRMS	zRMS Comments (if different PHI proposed)
Olive	14 days	Yes	14 days	
Grape	21 days	Yes	21 days	
Tomato, aubergine	10 days	Yes	10 days	
Pepper	14 days	n.a. (MRL exceedance)	-	Not recommended use
Leek, artichoke	7 days	No	-	Not recommended use
Celery, cardoon	7 days	No	-	Not recommended use
Flowering brassica	14 days	No	-	Not recommended use
Cucurbits edible peel	3 days	Yes	3 days	
Cucurbits inedible peel	7 days	n.a. (MRL exceedance)	-	Not recommended use
Onion	3 days	Yes	3 days	
Garlic, shallot	3 days	Yes	3 days	
Lettuce	7 days	No	-	Not recommended use
Other salad plants including leaves and sprouts of brassicacea	7 days	No	-	Not recommended use
Spinach and similar (leaves)	7 days	n.a. (MRL exceedance)	-	Not recommended use
Fresh beans/peas with pods	3 days	Yes	3 days	
Fresh beans without pod	3 days	Yes	3 days	
Fresh peas without pods, fresh lentils	3 days	No	-	Not recommended use
Potato	14 days	n.a. (MRL exceedance)	-	Not recommended use
Carrot	14 days	No	-	Not recommended use
Parsnip, Jerusalem artichoke, parsley, celeriac, horseradish and scorzonera	14 days	Yes	14 days	
Beetroot	14 days	No	-	Not recommended use
Strawberry	7 days	n.a. (MRL exceedance)	-	Not recommended use

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Crop	PHI for BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) requested by applicant	PHI/ Withholding period* sufficiently supported for Copper	PHI for BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) Blue proposed by zRMS	zRMS Comments (if different PHI proposed)
Ornamental plant and trees	NR		NR	Not assessed (non edible commodity)
Indoor uses				
Tomato, aubergine	3 days	Yes	3 days	
Pepper	14 days	n.a. (MRL exceedance)	-	Not recommended use
Cucurbits edible peel	3 days	Yes	3 days	
Cucurbits inedible peel	7 days	n.a. (MRL exceedance)	-	Not recommended use
Lettuce and other salad plants including leaves and sprouts of brassicacea	7 days	Yes	7 days	
Spinach and similar (leaves)	7 days	n.a. (MRL exceedance)	-	Not recommended use
Strawberry	7 days	n.a. (MRL exceedance)	-	Not recommended use
Ornamental plant and trees	NR		NR	Not assessed (non edible commodity)

NR: not relevant

* Purpose of withholding period to be specified

** F: PHI is defined by the application stage at last treatment (time elapsing between last treatment and harvest of the crop).

3.6 Environmental fate and behaviour (Part B, Section 8)

The fate and behaviour in the environment have been evaluated according to the requirements of Regulation (EC) No 1107/2009. Appropriate endpoints from the EU conclusions were used to calculate predicted environmental concentration (PEC) values for the active substance for the intended use patterns. In cases where deviations from the EU agreed endpoints were considered appropriate (for example when additional studies are provided), such deviations were highlighted and justified accordingly.

The PEC of copper in soil, surface water and groundwater have been assessed according to FOCUS guidance documents, with standard FOCUS scenarios to obtain outputs from the FOCUS models, and the endpoints established in the EU conclusions or agreed in the assessment based on new data provided.

No reliable PEC_{soil} were available for the active substance mainly due to a shorter period for estimating the accumulation in soil. Therefore, the risk assessment for the non-target terrestrial organisms cannot be finalised for all intended uses.

Given the uncertainties identified by zRMS in the notifier's exposure calculation (FOCUS STEP 1-2 for all entries to water bodies and FOCUS STEP 1-2 PEC_{sw} including mitigation measures) and the absence of results for all FOCUS scenarios, PEC_{sw} derived for the active substance cannot be used for the

ecotoxicological risk assessment. As a consequence, the risk assessment cannot be finalised for the non-target aquatic organisms.

PEC_{gw} for copper do not occur at levels exceeding those mentioned in Directive 98/83/CE¹⁴. Therefore, no unacceptable risk of groundwater contamination is expected for all intended uses.

Based on vapour pressure, no significant contamination of the air compartment is expected for the intended uses.

3.7 Ecotoxicology (Part B, Section 9)

The ecotoxicological risk assessment of the formulation was performed according to the requirements of Regulation (EC) No 1107/2009. Appropriate endpoints from the EU conclusions for the active substance were used for the intended use patterns. In cases where deviations from the EU agreed endpoints were considered appropriate (for example when additional studies are provided), such deviations were highlighted and justified accordingly.

An EFSA' Statement of the PPR panel on a framework for conducting the environmental exposure and risk assessment for transition metals when used as active substances in plant protection products was recently published (2021). This document provides useful recommendations upon applicability of new methodologies in the context of transition metals and possible areas of development for assessing the risk from transition metals used in PPPs. However, it does not provide valid tools for exposure assessment in the environment and toxicity estimation upon non-target organisms. Furthermore, no clear specific risk assessment schemes for transition metals used as active substances in PPPs is provided. Therefore, the risk assessment and conclusion are based on the methodology agreed by the experts during the renewal approval of the active substance. The EU-agreed endpoints recommended in the EFSA journal (EFSA Journal 2018;16(1):5152) were considered for the Art. 43 dossiers for copper compounds.

Based on the guidance documents, the risks for **non-target arthropods other than bee** and **non-target terrestrial plants** are acceptable for the intended uses.

For aquatic organisms, as the toxicity reference value for copper proposed by the applicant was based on an approach rejected at European level, it could not be used. In addition, no reliable PEC_{sd} were provided by the applicant for all uses. Therefore, the risk assessment for aquatic non-target species could not be finalised for uses in open field, in tunnels or in permanent greenhouses with soil-bound cultivation. For uses in permanent greenhouses with soil-less cultivation, the exposure of aquatic organisms to the active substance from the use of the product BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) is considered negligible.

For birds and mammals, the risk is not acceptable at Tier 1 for all intended uses. The arguments provided by the applicant to refine the risk assessment are identical to those that were considered insufficient at the European level. Therefore, without further data, the risk assessment for birds and mammals cannot be finalised except for applications under permanent greenhouse.

For bees, the risk assessment provided by the applicant is based on the EFSA Guidance Document¹⁵. For adult honey bees, the risk is not acceptable at Tier 1 for all intended uses. Higher-tier studies (cage and tunnel tests) are available and demonstrate that no adverse effects on adult honey bees are expected for all intended uses.

¹⁴ Council Directive 98/83/EC of 3 November 1998 on the quality of water intended for human consumption

¹⁵ EFSA Guidance Document on the risk assessment of plant protection products on bees (*Apis mellifera*, *Bombus* spp. and solitary bees) EFSA Journal 2013;11(7):3295

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

For honey bee larvae, the risks are not acceptable at Tier 1 for all intended uses except for tomatoes, lettuce, potatoes, olives, pistachios and hazelnuts in open-field or in walk-in tunnels. For all other uses in open-field or in walk-in tunnels, the higher-tier studies are not sufficient to demonstrate the absence of adverse effects of the product BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) on honey bee larvae. Therefore, the risk assessment for honey bee larvae cannot be finalised for applications in open-field or in walk-in tunnels except for tomatoes, lettuce, potatoes, olives, pistachios and hazelnuts.

For bumble bees, no acute risk assessment was provided by the applicant, although standard study protocols are available. Therefore, the risk assessment for bumble bees cannot be finalised for applications in open-field or in walk-in tunnels.

Overall, the risk for bees cannot be finalised for all requested uses except for applications under permanent greenhouse. For these structures, the following precautionary statement should be applied: “May affect pollinators. Avoid unnecessary exposure”.

For soil organisms, since no reliable PEC soil are available, a Tier 1 risk assessment cannot be conducted. For earthworms, the higher tier earthworm field trial data from a study conducted over 10 years with copper application every year demonstrates that there is an acceptable risk to earthworms for applications up to 4 kg Cu/ha/year. Therefore, an acceptable risk for earthworms is demonstrated for all intended uses of BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC).

For other soil meso- and macro-organisms, no higher-tier studies are available and extrapolating the results of the multiyear field study with earthworms to other soil meso- and macro-organisms was not supported by the experts at the Peer Review experts’ meeting 169. Therefore, the risk for soil macro-organisms other than earthworms could not be finalised for all intended uses except for uses under permanent greenhouse with soil-less cultivation.

For soil micro-organisms, based on a lack of effect at field level, the risks to soil micro-organisms are acceptable for the intended uses.

3.8 Relevance of metabolites (Part B, Section 10)

An assessment was conducted according to the SANCO/221/2000 guidance document. Please refer to environmental fate and behaviour above for conclusion on the risk of groundwater contamination.

4 Conclusion of the national comparative assessment (Art. 50 of Regulation (EC) No 1107/2009)

BOUILLIE BORDELAISE NC 20 K (BM 20 WP NC) contains copper compounds, which is approved as a candidate for substitution because it fulfills PBT criteria (Persistent and Toxic);

Steps 1 and 2 (French guidance document 27 July 2015):

- **Taking into account the agronomic interest, especially in the context of organic farming**

In accordance with Article 50, paragraphs 1.b) 1.c) and 1.d) of Regulation (EC) N°1107/2009,

- considering the absence of plant protection products or non-chemical methods of prevention or control allowing to consider a substitution of the product without major practical or economic disadvantage, and especially in the frame of organic farming,
- considering also the need to guarantee a diversity of the modes of action to reduce the emergence of resistance in target microorganisms,
- considering the need to take into account the minor uses of the product,

the substitution of the product will not be considered for all intended uses.

5 Further information to permit a decision to be made or to support a review of the conditions and restrictions associated with the authorisation

When the conclusions of the assessment is “Not acceptable”, please refer to relevant summary under point 3, “Background of authorisation decision and risk management”.

5.1.1 Post-authorisation monitoring

The monitoring of resistance to copper should implemented (one monitoring for all products based on copper) on *Xanthomonas sp.* on vegetable crops (e.g. tomato) and walnut and the results should be provided at the time of the next renewal of the product’s authorisation.

5.1.2 Post-authorisation data requirements

For renewal :

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Appendix 1 Copy of the product authorisation



BOUILLIEBO_PREX_2
019-4167_D.pdf

Copy of the product label

The draft product label as proposed by the applicant is reported below. The draft label may be corrected with consideration of any new element. The label shall reflect the detailed conditions stipulated in the Decision.

Mars 2019
Page 1 / 10

BOUILLIE BORDELAISE NC 20 K

FONGICIDE ET BACTÉRICIDE

Fongicide et bactéricide préventif à large spectre pour les vergers, la vigne, les cultures légumières, la pomme de terre, la betterave sucrière, les fraisiers et les cultures ornementales

Autorisation de Mise sur le Marché (A.M.M.) N° 2010509

Détenteur de l'A.M.M. : Industrias Químicas del Vallés, S.A. (Av. Rafael Casanova 81 - 08100 Mollet del Vallés - Barcelona, Espagne)

Contient 200 g/kg (20.0% p/p) de cuivre (sous forme de bouillie bordelaise)
Poudre mouillable (WP)

RESERVÉ À UN USAGE EXCLUSIVEMENT PROFESSIONNEL

BOUILLIE BORDELAISE NC 20 K (A.M.M. N° 2010509)		
Attention 	H410	Très toxique pour les organismes aquatiques, entraîne des effets néfastes à long terme.
	P391 P501	Recueillir le produit répandu. Éliminer le contenu/récipient conformément à la réglementation locale.
	Délai de rentrée sur la parcelle : 6 heures et, en cas d'application en milieu fermé, de 8 heures.	
	SPe3	Pour protéger les organismes aquatiques, respecter une zone non traitée par rapport aux points d'eau de : - 5 mètres avec dispositif végétalisé permanent de 5 mètres pour les usages sur agrumes, olive, vigne, tomate, aubergine, poivron, piment, poireau, oignon de printemps, ciboule, artichaut, céleri, cardon, brocoli, chou-fleur, concombre, courgette, cornichon, melon, pastèque, potiron, oignon, ail, échalote, laitue et autres salades, épinard, pourpier, feuilles de bettes, haricots non écosés et écosés frais, pois non écosés et écosés frais, lentilles fraîches, pomme de terre, betterave potagère, carotte et autres légumes racines, betterave sucrière et fraisier. - 20 mètres avec dispositif végétalisé permanent de 5 mètres pour les usages sur fruits à noyau, fruits à pépins et fruits à coque. - 20 mètres en plein champ pour les usages sur cultures ornementales.
	EUH401	Respectez les instructions d'utilisation afin d'éviter les risques pour la santé humaine et l'environnement.
SP1	Ne pas polluer l'eau avec le produit ou son emballage. Ne pas nettoyer le matériel d'application près des eaux de surface. Éviter la contamination via les systèmes d'évacuation des eaux à partir des cours de ferme ou des routes.	
FABRICANT :		
Adresse Tél. : xx-xx-xx-xx-xx		

EN CAS D'URGENCE

Composer le 15, le 112 ou contacter le centre anti-poison le plus proche

Puis signaler vos symptômes au réseau Phyt'Attitude, N° Vert : 0 800 887 887 (appel gratuit depuis un poste fixe).



Fiche de Données de Sécurité disponible sur demande pour les professionnels (ou sur www.quickfds.com).

Lire les instructions ci-jointes avant emploi.

EMB :

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Mars 2019
Page 2 / 10

N° de lot et date de fabrication : voir emballage
Version 1.0
XX kg e

PREMIERS SOINS

En cas d'inhalation :

Transporter la victime à l'air libre et la garder au chaud et au repos.

En cas de contact avec les yeux :

Laver abondamment avec de l'eau douce et propre durant au minimum 15 minutes en maintenant les paupières écartées. Retirer les lentilles de contact si la victime en porte après 5 minutes de rinçage. Consulter un spécialiste.

En cas de contact avec la peau :

Enlever les vêtements imprégnés et laver soigneusement la peau avec de l'eau et du savon ou utiliser un nettoyant connu. Ne pas utiliser de solvants ou de diluants. En cas d'irritation, consulter un médecin.

En cas d'ingestion :

Rincer la bouche avec de l'eau et consulter un médecin ou un centre ~~anti-poison~~ ^{antipoison}. Montrer l'étiquette. Garder au repos.

Ne pas faire vomir.

Antidote : EDTA, BAL ou ~~pénicillamine~~ ^{pénicillamine}. Consulter immédiatement un médecin et lui montrer l'étiquette.

Traitement symptomatique : un lavage gastrique avec une solution ~~lacté~~ ^{lacté}-albumine peut être nécessaire.

RECOMMANDATIONS D'EMPLOI

IMPORTANT : Lire attentivement les instructions de cette section afin de garantir une utilisation sûre et efficace de ce produit.

MODE D'ACTION

Le cuivre appartient au groupe FRAC M01. C'est un fongicide et bactéricide multi-site inorganique.

TABLEAU DES USAGES

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Citron, Mandarine, Orange, Pamplemousse, Clémentine, Limette	Alternariose (<i>Alternaria alternata</i> f. sp. <i>Citri</i>) Bactérioses (<i>Pseudomonas syringae</i> pv. <i>syringae</i>) Chancres du collet (<i>Phytophthora</i> spp.)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (feuilles visibles mais n'ont pas encore atteint leur taille finale) à 89 (fruits mûrs)	1000-2000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Abricotier, Cerisier, Pêcher, Nectarinier, Prunier, Mirabellier, Jujubier	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>syringae</i>) Chancres à champignons (<i>Cytospora leucostoma</i>) Cloques (<i>Taphrina deformans</i>) Cancroème (<i>Stigmella camponotile</i>) Monilioses (<i>Monilia</i> spp.)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 95 (sénescence - 50% des feuilles sont décolorées ou tombées) à 53 (éclatement des bourgeons)	400-1000 L/ha	-	20 mètres avec DVP** de 5 mètres
Cerisier	Anthraxose (<i>Blumeriella jappa</i>)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 95 (50% des feuilles sont décolorées ou tombées) à 53 (éclatement des bourgeons)	400-1000 L/ha	-	20 mètres avec DVP** de 5 mètres
Pommier, Poirier, Cognassier, Néflier, Nashi, Pommette	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>populans</i>) Chancres européens (<i>Nectria galligena</i>) Feu bactérien (<i>Erwinia amylovora</i>) Monilioses (<i>Monilia</i> spp.) Tavelure (<i>Venturia</i> spp.)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 91 (début sénescence - fin de la croissance des rameaux, le bourgeon terminal est développé, les feuilles sont toujours vertes) à 53 (éclatement des bourgeons)	500-1000 L/ha	-	20 mètres avec DVP** de 5 mètres

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Mars 2019
Page 4 / 10

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Amandier	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>syringae</i>) Cloques (<i>Taphrina deformans</i>) Corneum (<i>Stigmella carpophila</i>) Monilioses (<i>Monilia</i> spp.)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 95 (50% des feuilles sont décolorées ou tombées) à 53 (éclatement des bourgeons)	500-1000 L/ha	-	20 mètres avec DVP** de 5 mètres
Noyer, Noisetier, Châtaignier, Pistachier	Alternariose (<i>Alternaria</i> spp.) Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Dépérissement cryptogamique (<i>Cytospora</i> spp.) Sépiose (<i>Mycosphaerella</i> spp.)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 95 (50% des feuilles sont décolorées ou tombées) à 53 (éclatement des bourgeons)	500-1000 L/ha	-	20 mètres avec DVP** de 5 mètres
Olivier	Anthraxose (<i>Colletotrichum gloeosporioides</i>) Bactérioses (<i>Pseudomonas savastanoi</i> pv. <i>savastanoi</i>) Maladie de l'œil de paon (<i>Spilocaea oleaginea</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 10 (développement des feuilles) à 85 (début de la coloration du fruit)	800-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Vigne (raisin de cuve et raisin de table) (champ)	Anthraxose (<i>Elsinoe ampelina</i>) Bactérioses (<i>Xanthomonas ampelina</i>) Mildiou (<i>Plasmopara viticola</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (3 feuilles étalées) à 81 (début de la maturation) et 91 (après la vendange : aoûté du bois terminé)	400-1000 L/ha	21 jours	5 mètres avec DVP** de 5 mètres
Tomate, Aubergine (champ/ sous serre)	Bactérioses (<i>Pseudomonas syringae</i> et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria solani</i> , <i>Colletotrichum</i> spp.) Mildiou (<i>Phytophthora infestans</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (5 feuilles étalées sur la tige principale) à 89 (fruits mûrs)	400-1000 L/ha	10 jours (plein champ) / 3 jours (sous serre)	5 mètres avec DVP** de 5 mètres
Poivron, Piment (champ/ sous serre)	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>Syringae</i> et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria solani</i> , <i>Colletotrichum</i> spp.) Mildiou (<i>Phytophthora capsici</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (5 feuilles étalées sur la tige principale) à 89 (fruits mûrs)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Poireau, Oignon de printemps, Ciboule Artichaut, Céleri (branche et rave), Cardon (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria</i> spp., <i>Ascochyta</i> spp.) Mildiou (<i>Bremia lactucae</i> , <i>Phytophthora porri</i> , <i>Plasmopara crustacea</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 14 (4 feuilles visibles/étalées) à 51 (le bulbe commence à s'allonger/ la pousse principale commence à sortir)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Mars 2019
Page 5 / 10

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Brocoli, Chou-fleur (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas campestris</i> pv. <i>campestris</i>) Maladies des taches brunes (<i>Alternaria brassicae</i> , <i>Colletotrichum</i> spp.) Mildiou (<i>Hyaloperonospora brassicae</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 14 (4 feuilles étalées) à 59 (premiers pétales visibles, fleurs toujours fermées)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Concombre, Courgette, Cornichon (champ/ sous serre)	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>Lachrymans</i> et <i>Xanthomonas campestris</i> pc. <i>cucurbitae</i>) Maladies des taches brunes (<i>Alternaria cucumerina</i> , <i>Colletotrichum lagenarium</i>) Mildiou (<i>Pseudoperonospora cubensis</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 10 (cotylédons étalés) à 89 (maturation complète)	400-1000 L/ha	3 jours	5 mètres avec DVP** de 5 mètres
Melon, Pastèque, Potiron (champ/ sous serre)	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>Lachrymans</i> et <i>Xanthomonas campestris</i> pc. <i>cucurbitae</i>) Maladies des taches brunes (<i>Alternaria cucumerina</i> , <i>Colletotrichum lagenarium</i>) Mildiou (<i>Pseudoperonospora cubensis</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 10 (cotylédons étalés) à 89 (maturation complète)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres
Oignon, Ail, Echalote (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria porri</i> , <i>Colletotrichum coconodes</i> , <i>Pleospora alli</i>) Mildiou (<i>Peronospora destructor</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 14 (4 feuilles visibles) à 47 (feuilles fanées dans 10% des plantes)	400-1000 L/ha	3 jours	5 mètres avec DVP** de 5 mètres
Laitue et autres salades : Mâche Scarole (endive à larges feuilles) Cresson Cresson de terre Rucola, Rucola Moutarde brune Feuilles et pousses de Brassica spp. (champ/ sous serre)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas axonopodis</i> pv. <i>Vitians</i>) Mildiou (<i>Bremia lactucae</i>) Maladies des taches brunes (<i>Alternaria sonchi</i> , <i>Marssonina panattoniana</i>) Sclerotinioses (<i>Sclerotinia minor</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 12 (2 feuilles étalées) à 49 (la taille typique de la masse foliaire est atteinte)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres
Epinard, Pourpier, Feuilles de bettes (champ/ sous serre)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria</i> spp., <i>Colletotrichum spinaciae</i>) Mildiou (<i>Peronospora farinosa</i> f. sp. <i>Spinaciae</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 12 (2 feuilles étalées) à 49 (la taille typique de la masse foliaire est atteinte)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres

BM 20 WP NC / BOUILLIE BORDELAISE NC 20 K
Part A - National Assessment
FRANCE

Mars 2019
Page 6 / 10

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Haricots non écosés frais (haricots verts, etc.) Haricots écosés frais (fèves, flageolets, etc.) Pois non écosés frais (pois mange-tout) Pois écosés frais (pois potagers, pois frais, pois chiches) Lentilles fraîches (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas axonopodis</i> pv. <i>phaseoli</i>) Maladies des taches brunes (<i>Colletotrichum lindemuthianum</i> , <i>Sclerotinia</i> spp.) Mildiou (<i>Peronospora</i> viciae, <i>Phytophthora phaseoli</i>) Rouilles (<i>Uromyces</i> spp.)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 14 (4 feuilles étalées ou 4 vrilles formées) à 89 (maturation complète)	400-1000 L/ha	3 jours	5 mètres avec DVP** de 5 mètres
Pomme de terre	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria solani</i> , <i>Colletotrichum coccodes</i>) Mildiou (<i>Phytophthora infestans</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (5*** feuille de la tige principale étalée) à 85 (baies de la première inflorescence ocre ou brunâtres)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Betterave potagère Carotte Raifort Topinambours Panais Persil à grosse racine Radis Salsifis Rutabagas Navets (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas campestris</i> pv. <i>carotae</i>) Maladies des taches brunes (<i>Alternaria dauci</i> , <i>Cercospora carotae</i> , <i>Colletotrichum</i> spp.) Mildiou (<i>Plasmopara nivea</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (5 feuilles étalées) à 47 (racines, tiges ou tubercules ont atteint 70% de leur taille finale)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Betterave sucrière	Maladies du feuillage (<i>Cercospora beticola</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 39 (fermeture des lignes: environ 90% des plantes des lignes adjacentes se touchent) à 49 (racine de la betterave à sa taille de récolte)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Fraisier (champ/ sous serre)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas fragariae</i>) Maladies des taches brunes (<i>Colletotrichum fragariae</i> , <i>Mycosphaerella fragariae</i>) Mildiou (<i>Phytophthora fragariae</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 13 (3 feuilles étalées) à 85 (les premières fraises ont atteint la couleur spécifique de la variété)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Arbres et Arbustes d'ornement Bulbes ornementaux Cultures florales et plantes vertes Rosier (champ/ sous serre)	Anthracnoses et maladies des taches foliaires (<i>Colletotrichum spp.</i> , <i>Septoria spp.</i>) Mildiou (<i>Peronospora spp.</i>) Monilioses (<i>Monilia spp.</i>) Rouilles (<i>Puccinia spp.</i>)	3.75 à 5 kg/ha	3 (intervalle de 7 jours entre appl.)	Tous les stades	400-1000 L/ha	-	20 mètres en plein champ

* Seules les utilisations entraînant une application totale maximale de 28 kg de cuivre par hectare sur une période de 7 ans sont autorisées.

** DVP : Dispositif Végétalisé Permanent

LIMITES MAXIMALES DE RÉSIDUS (LMR)

Les LMR sont consultables à l'adresse suivante : <http://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/>

PRÉCONISATIONS D'EMPLOI

BOUILLIE BORDELAISE NC 20 K est un fongicide et bactéricide préventif à large spectre. A base de cuivre, il est plus efficace contre les spores que contre les maladies déjà établies, il est donc recommandé de l'appliquer avant l'apparition de la maladie, ou dès que les premiers symptômes de la maladie sont visibles.

Dans des conditions très favorables au développement des maladies visées, il est recommandé d'alterner l'utilisation de BOUILLIE BORDELAISE NC 20 K avec d'autres produits contenant des substances actives ayant un mode d'action différent.

PRÉPARATION DE LA BOUILLIE

Remplir à moitié la cuve avec de l'eau et mettre en marche l'agitation. Verser la quantité nécessaire de BOUILLIE BORDELAISE NC 20 K dans la cuve du pulvérisateur. Remplir la cuve avec de l'eau au volume requis. Maintenir l'agitation durant toute la durée de l'application.

APPLICATION

Éviter tout risque de dérive de pulvérisation. Traiter par temps calme.

Ne pas laisser la bouillie dans la cuve du pulvérisateur pendant de longues périodes par exemple le temps des repas.

Appliquer en pulvérisation foliaire. Pour assurer une efficacité optimale, le matériel de pulvérisation doit être calibré afin de bien appliquer le produit sur l'ensemble des parties de la culture à protéger et assurer une distribution uniforme du produit sur la culture. Ajuster le volume de bouillie en fonction du matériel utilisé et du volume de végétation afin d'assurer une couverture complète de la végétation à protéger.

Appliquer préventivement avec un intervalle recommandé de 7 jours entre les applications. Cependant, en règle générale, les applications doivent toujours être adaptées à l'évolution de la maladie, de la culture traitée et aux conditions climatiques.

MÉLANGES

Les mélanges extemporanés doivent être mis en œuvre conformément à la réglementation en vigueur et aux recommandations des guides de bonnes pratiques officiels.

BOUILLIE BORDELAISE NC 20 K est compatible avec la plupart des produits phytopharmaceutiques couramment utilisés, à l'exception de ceux ayant une forte réaction acide ou alcaline. Pour les mélanges, il est donc recommandé d'effectuer un test préalable avec une petite quantité de produits avant de généraliser le mélange.

Protection de l'opérateur et du travailleur

Se laver les mains après toute manipulation/utilisation/intervention dans une parcelle préalablement traitée.

Ne pas manger, boire, téléphoner ou fumer lors de l'utilisation du produit.

Caractéristiques des EPI ▼		PROTECTION DE L'UTILISATEUR PENDANT LES PHASES DE :			PROTECTION DU TRAVAILLEUR en cas d'intervention sur les parcelles traitées
		MÉLANGE/ CHARGEMENT et NETTOYAGE	APPLICATION AVEC PULVÉRISATEUR PORTÉ OU TRAINÉ À RAMPE, PNEUMATIQUE OU ATOMISEUR TRACTEUR AVEC OU SANS CABINE	APPLICATION MANUELLE LANCE OU PULVÉRISATEUR À DOS	
CANTS EN NITRILE réutilisables (certifiés EN 374-3) ou à usage unique (certifiés EN 374-2)		Réutilisables	À usage unique, si intervention sur le matériel de pulvérisation	Réutilisables	
EPI VESTIMENTAIRE 65% polyester / 35% coton ≥ 230 g/m² + traitement déperlant		EPI vestimentaire ET EPI partiel			
EPI PARTIEL blouse ou tablier à manches longues catégorie III type PB3 certifié EN14605+A1					
LUNETTES ou ECRAN FACIAL certifiés EN 166:2002 (CE, sigle 3)					
PROTECTION RESPIRATOIRE masque (EN 136:1998 ou EN 140:1998) équipé d'un filtre P3 (EN143:2006) minimum					
BOTTES certifiées EN 13 832-3:2006					

Pour protéger l'opérateur :

- pendant le mélange/chargement et le nettoyage du matériel de pulvérisation, porter :

- Gants en nitrile certifiés EN 374-3,
- Combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement déperlant.
- EPI partiel (blouse ou tablier à manches longues) de catégorie III et de type PB (3) à porter par-dessus la combinaison précitée,
- Lunettes ou écran facial certifiés norme EN 166 (CE, sigle 3),
- Protection respiratoire certifiée : conforme aux normes EN 136 (masque complet) ou EN 140 (demi-masque) et équipée d'un filtre P3 (EN 143) au minimum.

- pendant l'application, porter :

Application avec tracteur avec cabine :

- Combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement déperlant.
- Gants en nitrile certifiés EN 374-2 à usage unique dans le cas d'une intervention sur le matériel pendant la phase de pulvérisation. Dans ce cas, les gants ne doivent être portés qu'à l'extérieur de la cabine et doivent être stockés après utilisation à l'extérieur de la cabine.

Application avec tracteur sans cabine :

- Combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement déperlant.
- Gants en nitrile certifiés EN 374-2 à usage unique, dans le cas d'une intervention sur le matériel pendant la phase de pulvérisation.

Application manuelle avec une lance ou un pulvérisateur à dos :

- Combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement ~~déperlant~~.
- Gants en nitrile certifiés EN 374-3,
- Bottes de protection certifiées EN 13 832-3.

Pour protéger le travailleur :

Dans les cas où le travailleur serait amené à intervenir sur les parcelles traitées, porter une combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement ~~déperlant~~.

Rapporter les équipements de protection individuelle (EPI) usagés dans un sac translucide à votre distributeur partenaire ECO EPI ou faire appel à une entreprise habilitée pour la collecte et l'élimination de produits dangereux.

RÉSISTANCE

L'utilisation répétée, sur une même parcelle, de préparations à base de substances actives de la même famille chimique ou ayant le même mode d'action, peut conduire à l'apparition d'organismes résistants.

Pour réduire ce risque, l'utilisateur doit raisonner en premier lieu les pratiques agronomiques et respecter les conditions d'emploi du produit.

Il est conseillé d'alterner ou d'associer, sur une même parcelle, des préparations à base de substances actives de familles chimiques différentes ou à modes d'action différents, tant au cours d'une saison culturale que dans la rotation.

En dépit du respect de ces règles, on ne peut pas exclure une altération de l'efficacité de cette préparation liée à ces phénomènes de résistance. De ce fait, Industrias Químicas del Vallés décline toute responsabilité quant à d'éventuelles conséquences qui pourraient être dues à de telles résistances.

STOCKAGE

Conserver le produit uniquement dans son emballage d'origine, dans un local phytopharmaceutique conforme à la réglementation en vigueur, à l'écart des aliments et boissons, y compris ceux pour animaux.

Conserver hors de la portée des enfants et des personnes non autorisées.

NETTOYAGE DU PULVÉRISATEUR ET GESTION DES FONDS DE CUVE

À la fin de la période d'application avec le produit, l'intégralité de l'appareil (cuve, rampe, circuit, buses...) doit être rincée à l'eau claire additionnée d'un mouillant (recommandé pour le nettoyage des pulvérisateurs). Le rinçage du pulvérisateur, l'épandage ou la vidange du fond de cuve et l'élimination des effluents doivent être réalisés conformément à la réglementation en vigueur.

ÉLIMINATION DU PRODUIT ET DE SON EMBALLAGE

Réemploi de l'emballage interdit.

Apporter les emballages vidés et pliés à votre distributeur partenaire d'A.D.I.VALOR ou à un autre service de collecte spécifique.

Pour l'élimination des produits non utilisables, conserver le produit dans son emballage d'origine. Interroger votre distributeur partenaire d'A.D.I.VALOR ou faites appel à une entreprise habilitée pour la collecte et l'élimination des déchets dangereux.

En cas de déversement accidentel :

Se protéger (EPI) et sécuriser la zone. Prévenir les pompiers (18 ou 112) en cas de danger immédiat pour l'environnement que vous ne pouvez gérer avec vos propres moyens.

Collecter tout ce qui a pu être en contact avec le produit, terre souillée incluse. Nettoyer le site et le matériel utilisé, en prenant soin de confiner les effluents générés par l'opération de nettoyage. Les éliminer selon la réglementation en vigueur.



AVERTISSEMENT

Toute reproduction totale ou partielle de cette étiquette est interdite.

Respecter les usages, doses, conditions et précautions d'emploi mentionnés sur l'emballage. Ils ont été déterminés en fonction des caractéristiques du produit et des applications pour lesquelles il est préconisé. Conduire sur ces bases la culture et les traitements selon la bonne pratique agricole en tenant compte, sous la responsabilité de l'utilisateur, de tous les facteurs particuliers concernant votre exploitation, tels que la nature du sol, les conditions météorologiques, les méthodes culturales, les variétés végétales, la résistance des espèces...

Le fabricant garantit la qualité du produit vendu dans son emballage d'origine et stocké selon les conditions préconisées, ainsi que sa conformité à l'Autorisation de Mise sur le Marché délivrée par les autorités compétentes françaises. Pour les denrées issues de cultures protégées avec cette spécialité et destinées à l'exportation, il est de la responsabilité de l'exportateur de s'assurer de la conformité avec la réglementation en vigueur dans le pays importateur.