

REGISTRATION REPORT

Part A

Risk Management

Product code: Super Bouillie Macclesfield 80

Product name: SUPER BOUILLIE MACCLESFIELD 80

Chemical active substance:

copper, 200 g/kg

Southern Zone & Interzonal

Zonal Rapporteur Member State: France

NATIONAL ASSESSMENT FRANCE

(authorisation renewal)

Applicant: Industrias Químicas del Vallés S.A. (IQV)

Date: 15/07/2025

Table of Contents

1	Details of the application	4
1.1	Application background	4
1.2	Letters of Access.....	5
1.3	Justification for submission of tests and studies.....	5
1.4	Data protection claims.....	5
2	Details of the authorisation decision.....	6
2.1	Product identity	6
2.2	Conclusion DAMM	6
2.3	Substances of concern for national monitoring	6
2.4	Classification and labelling.....	6
2.4.1	Classification and labelling under Regulation (EC) No 1272/2008	6
2.4.2	Standard phrases under Regulation (EU) No 547/2011	7
2.4.3	Other phrases (according to Article 65 (3) of the Regulation (EU) No 1107/2009)	7
2.5	Risk management	7
2.5.1	Restrictions linked to the PPP	8
2.5.2	Specific restrictions linked to the intended uses.....	8
2.6	Intended uses (only NATIONAL GAP)	9
3	Background of authorisation decision and risk management	53
3.1	Physical and chemical properties (Part B, Section 2)	53
3.2	Efficacy (Part B, Section 3)	53
3.3	Methods of analysis (Part B, Section 5)	53
3.3.1	Analytical method for the formulation.....	53
3.3.2	Analytical methods for residues	53
3.4	Mammalian toxicology (Part B, Section 6)	54
3.4.1	Acute toxicity.....	54
3.4.2	Operator exposure.....	54
3.4.3	Worker exposure.....	57
3.4.4	Bystander exposure	61
3.4.5	Resident exposure.....	61
3.4.6	Combined exposure.....	63
3.5	Residues and consumer exposure (Part B, Section 7)	63
3.6	Environmental fate and behaviour (Part B, Section 8).....	65
3.7	Ecotoxicology (Part B, Section 9)	66
3.8	Relevance of metabolites (Part B, Section 10)	67
4	Conclusion of the national comparative assessment (Art. 50 of Regulation (EC) No 1107/2009)	67

5	Further information to permit a decision to be made or to support a review of the conditions and restrictions associated with the authorisation.....	68
5.1.1	Post-authorisation monitoring.....	68
5.1.2	Post-authorisation data requirements	68
Appendix 1	Copy of the product authorisation DAMM	69
Appendix 2	Copy of the product label	86

PART A

RISK MANAGEMENT

1 Details of the application

The company INDUSTRIAS QUÍMICAS DEL VALLÉS S.A. has requested a marketing authorisation in France for the product SUPER BOUILLIE MACCLESFIELD 80 (formulation code: Super Bouillie Macclesfield 80), containing 200 g/kg copper¹ (in the form of Bordeaux mixture (CAS n° 8011-63-0)) as a fungicide and bactericide for professional uses.

Appendix 1 of this document provides a copy of the product authorisation.

Appendix 2 of this document contains a copy of the product label (draft as proposed by the applicant).

Appendix 3 of this document is the list of data considered for national authorisation.

1.1 Application background

The present registration report concerns the evaluation of INDUSTRIAS QUÍMICAS DEL VALLÉS S.A.'s application submitted on 01/04/2019 to market SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) in France (product uses described under point 2.3). France acted as a zonal Rapporteur Member State (zRMS) for field uses and interzonal Rapporteur Member State (izRMS) for greenhouse uses for this request and assessed the application submitted for the re-registration of authorisation after the renewal of approval of the active substances copper compounds of this product in France and in other Member States (MSs) of the European Union.

The present application (2019-3931) was evaluated in France by the French Agency for Food, Environmental and Occupational Health & Safety (Anses), according to the Regulation (EC) n° 1107/2009², the implementing regulations, and French regulations. This application was assessed in the context of the zonal procedure for all MSs of the Southern zone for field uses and for all MSs of the European Union for greenhouse uses, taking into account the worst-case uses ("risk envelope approach")³. When risk mitigation measures were necessary, they are adapted to the situation in France.

The data taken into account are those deemed to be valid either at European level (Review Report and EFSA conclusion) or at zonal/national level. The assessment of SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) has been made using endpoints agreed in the EU peer review of copper compounds. It also includes assessment of data and information related to SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) where those data have not been considered in the EU peer review process.

¹ COMMISSION IMPLEMENTING REGULATION (EU) 2018/1981 of 13 December 2018 renewing the approval of the active substances copper compounds, as candidates for substitution, in accordance with Regulation (EC) No 1107/2009 of the European Parliament and of the Council concerning the placing of plant protection products on the market, and amending the Annex to Commission Implementing Regulation (EU) No 540/2011

² REGULATION (EC) No 1107/2009 of the European Parliament and of the Council of 21 October 2009 concerning the placing of plant protection products on the market and repealing Council Directives 79/117/EEC and 91/414/EEC

³ SANCO document "risk envelope approach", European Commission (14 March 2011). [Guidance document on the preparation and submission of dossiers for plant protection products according to the "risk envelope approach"; SANCO/11244/2011 rev. 5](#)

The conclusions of the assessment published by EFSA 2018^{4,5}, as part of the procedure for the renewal of the approval of copper compounds, based on the available information, identify risk for non-target organisms for the representative uses on grapevine, cucurbits and tomatoes, as well as to workers for the grapevine use.

In the framework of MRL review for copper compounds under Article 12 of Regulation (CE) 396/2005, EFSA published a reasoned opinion (EFSA, 2018⁶). Based on an evaluation of the available data MRL have been proposed and a consumer risk assessment has been conducted. Some information required by the regulation has not been transmitted and a chronic risk for the consumers was identified. Therefore the consumer risk assessment is only tentative and some of the proposed MRL still require a decision by risk managers. Exposure reduction measures could also be investigated.

This part A of the RR presents a summary of essential scientific points upon which recommendations are based and is not intended to show the assessment in detail. The risk assessment conclusions provided in this document are based on the information, data and assessments provided in the Registration Report, Part B Sections 1-10 and Part C, and where appropriate the addendum for France.

The conclusions on the acceptability of risk are based on the criteria provided in Regulation (EU) No 546/2011⁷, and are expressed as “acceptable” or “not acceptable” in accordance with those criteria.

This document also describes the specific conditions of use and labelling required for France for the registration of SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80).

1.2 Letters of Access

The applicant is the owner of data which support the renewal of approval of the active substance. The applicant has provided letters of access for active substance and PPP data. These letters of access are available upon request.

1.3 Justification for submission of tests and studies

According to the applicant: *“The new studies afforded in this dossier, not previously assessed, provided in this dossier are necessary according to the requirements and guidelines into force which didn’t exist when the product was assessed by Uniform Principles in the past.”*

1.4 Data protection claims

Data protection is claimed in accordance with Article 59 of Regulation (EC) No. 1107/2009 as provided for in the list of references in Appendix 3.

⁴ Peer review of the pesticide risk assessment of the active substance copper compounds Copper(I), copper(II) variants namely copper hydroxide, copper oxychloride, tribasic copper sulfate, copper(I) oxide, Bordeaux mixture, EFSA Journal 2018;16(1):515

⁵ Outcome of the consultation with Member States, the applicant and EFSA on the pesticide risk assessment for copper compounds copper(I), copper(II) variants namely copper hydroxide, copper oxychloride, tribasic copper sulfate, copper(I) oxide, Bordeaux mixture in light of confirmatory data. EFSA supporting publication 2018:EN-1486.

⁶ REASONED OPINION ADOPTED: 1 March 2018. Review of the existing maximum residue levels for copper compounds according to Article 12 of Regulation (EC) No 396/2005 European Food Safety Authority (EFSA).

⁷ COMMISSION REGULATION (EU) No 546/2011 of 10 June 2011 implementing Regulation (EC) No 1107/2009 of the European Parliament and of the Council as regards uniform principles for evaluation and authorisation of plant protection products

2 Details of the authorisation decision

2.1 Product identity

Product code	Super Bouillie Macclesfield 80
Product name in MS	SUPER BOUILLE MACCLESFIELD 80
Authorisation number	7500716
Kind of use	Professional use
Low risk product (article 47)	No
Function	Fungicide and bactericide
Applicant	INDUSTRIAS QUÍMICAS DEL VALLÉS S.A.
Active substance(s) (incl. content)	copper, 200 g/kg (in the form of Bordeaux mixture)
Formulation type	Wettable powder [WP]
Packaging	PET/LDPE bag (1 kg, 5 kg) paper/LDPE bag (5 kg, 10 kg, 20 kg) OPP/PET met/LDPE bag (1 kg, 5 kg) PP big bag (1000 kg)
Coformulants of concern for national authorisations	-
Restrictions related to identity	-
Mandatory tank mixtures	None
Recommended tank mixtures	None

2.2 Conclusion DAMM

The evaluation of the application for SUPER BOUILLE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) resulted in the decision **to refuse** the authorisation.

2.3 Substances of concern for national monitoring

Refer to 5.1.1.

2.4 Classification and labelling

2.4.1 Classification and labelling under Regulation (EC) No 1272/2008

The following classification is proposed in accordance with Regulation (EC) No 1272/2008:

Hazard class(es), categories:	Acute toxicity (inhalation), category 4 Hazardous to the aquatic environment - Acute Hazard, category 1 Hazardous to the aquatic environment - Chronic Hazard, category 1
-------------------------------	---

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Hazard pictograms:	  GHS07 GHS09
Signal word:	Warning
Hazard statement(s):	H332: Harmful if inhaled. H400: Very toxic to aquatic life. H410: Very toxic to aquatic life with long-lasting effects.
Precautionary statement(s):	<i>For the P phrases, refer to the existing legislation</i>
Additional labelling phrases:	None

See Part C for justifications of the classification and labelling proposals.

2.4.2 Standard phrases under Regulation (EU) No 547/2011

NA

2.4.3 Other phrases (according to Article 65 (3) of the Regulation (EU) No 1107/2009)

None.

2.5 Risk management

According to the French law and procedures, specific conditions of use are set out in the Decision letter. The French Order of 4 May 2017⁸ provides that:

- unless otherwise stated in the product authorisation, the pre harvest interval (PHI) is at least 3 days;
- unless otherwise stated in the product authorisation, the minimum buffer zone alongside a water body is 5 metres for products applied through spraying or dusting;
- unless otherwise stated in the product authorisation, the minimum re-entry period is 6 hours for field uses and 8 hours for indoor uses.

Drift reduction measures such as low-drift nozzles are not considered within the decision-making process in France. However, non-spraying buffer zones may be reduced under some circumstances as explained in appendix 3 of the above-mentioned French Order.

Finally, the French Order of 26 March 2014⁹ provides that:

- an authorisation granted for a “reference” crop applies also for “related” crops, unless formally stated in the Decision
- the “reference” and “related” crops are defined in Appendix 1 of that French Order.

Thus, at French national level, possible extrapolation of submitted data and the corresponding assessment from “reference” crops to “related” ones are undertaken even if not clearly requested by the applicant in their dRR, and a conclusion is also reached on the acceptability of the intended uses on those “related”

⁸ Arrêté du 4 mai 2017 relatif à la mise sur le marché et à l'utilisation des produits phytopharmaceutiques et de leurs adjuvants visés à l'article L. 253-1 du code rural et de la pêche maritime, modifié par l'arrêté du 27 décembre 2019 <https://www.legifrance.gouv.fr/eli/arrete/2017/5/4/AGRGI632554A/jo/texte> ; <https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000039686039&categorieLien=id>

⁹ <http://www.legifrance.gouv.fr/eli/arrete/2014/3/26/AGRGI407093A/jo>

crops. The aim of this Order, mainly based on the EU document on residue data extrapolation¹⁰ is to supply “minor” crops with registered plant protection products.

Therefore the GAP table (Section 2.3) and Decision may include uses on crops not originally requested by the applicant.

The Decision, as reproduced in Appendix 1, takes also into account national provisions, including national mitigation measures.

2.5.1 Restrictions linked to the PPP

NA :

2.5.2 Specific restrictions linked to the intended uses

Some of the authorised uses are linked to the following conditions in addition to those listed under point 2.5.1 (mandatory labelling):

None.

¹⁰ SANCO document “guidance document:- Guidelines on comparability, extrapolation, group tolerances and data requirements for setting MRLs”: SANCO/ 7525/VI/95 - rev.9

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

2.6 Intended uses (only NATIONAL GAP)

Please note: The GAP Table below reports the intended uses proposed by the applicant, and possible extrapolation according to French Order of 26 March 2014 (highlighted in green), evaluated and concluded as safe uses by France as zRMS (field uses) and izRMS (greenhouse uses). Those uses are then granted in France.

When the conclusion is “not acceptable” or “not finalised”, the intended use is highlighted in grey and the main reason(s) reported in the remarks.

When a use is “acceptable” with GAP restrictions, the modifications of the GAP are in bold.

Use should be crossed out when the applicant no longer supports this use.

GAP rev. 1, date: 2025-07-15

PPP (product name/code): SUPER BOUILLIE MACCLESFIELD 80
Active substance 1: copper (Bordeaux mixture)
Applicant: INDUSTRIAS QUÍMICAS DEL VALLÉS S.A.
Zone(s): Southern Zone & Interzonal ^(d)
Verified by MS: Yes
Field of use: Fungicide and bactericide

Formulation type: WP ^(a, b)
Conc. of a.s. 1: 200 g/kg ^(c)
Professional use: ☒
Non-professional use: ☐

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
Zonal uses (field or outdoor uses, certain types of protected crops)													
1	FR (SEZ)	Lemon <i>(Citrus limon</i> ; 0110030; FC 0002, FC 0204)	F	Phytophthora <i>(Phytophthora spp.</i> ; 1PHYTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	Not acceptable (worker, PPE*)
2	FR (SEZ)	Lemon <i>(Citrus limon</i> ; 0110030; FC 0002, FC 0204)	F	Blast or black pit <i>(Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
3	FR (SEZ)	Lemon <i>(Citrus limon;</i> 0110030; FC 0002, FC 0204)	F	Brown spot <i>(Alternaria alternata f. sp.</i> <i>citri</i> ; ALTEAC)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
4	FR (SEZ)	Mandarin <i>(Citrus reticulata;</i> 0110050; FC 0003, FC 0206)	F	Phytophthora <i>(Phytophthora spp.;</i> 1PHYTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	Not acceptable (worker, PPE*)
5	FR (SEZ)	Mandarin <i>(Citrus reticulata;</i> 0110050; FC 0003, FC 0206)	F	Blast or black pit <i>(Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
6	FR (SEZ)	Mandarin <i>(Citrus reticulata;</i> 0110050; FC 0003, FC 0206)	F	Brown spot <i>(Alternaria alternata f. sp.</i> <i>citri</i> ; ALTEAC)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
7	FR (SEZ)	Orange <i>(Citrus sp;</i> 0110020; FC 0004, FC 0208)	F	Phytophthora <i>(Phytophthora spp.;</i> 1PHYTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
8	FR (SEZ)	Orange <i>(Citrus sp;</i> 0110020; FC 0004, FC 0208)	F	Blast or black pit <i>(Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	Not acceptable (worker, PPE*)
9	FR (SEZ)	Orange <i>(Citrus sp;</i> 0110020; FC 0004, FC 0208)	F	Brown spot <i>(Alternaria alternata f. sp.</i> <i>citri</i> ; ALTEAC)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
10	FR (SEZ)	Grapefruit <i>(Citrus paradise;</i> 0110010; FC 0005, FC 0203)	F	Phytophthora <i>(Phytophthora spp.;</i> 1PHYTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	Not acceptable (worker, PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
11	FR (SEZ)	Grapefruit (<i>Citrus paradise</i> ; 0110010; FC 0005, FC 0203)	F	Blast or black pit (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
12	FR (SEZ)	Grapefruit (<i>Citrus paradise</i> ; 0110010; FC 0005, FC 0203)	F	Brown spot (<i>Alternaria alternata</i> f. sp. <i>citri</i> ; ALTEAC)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	1000- 2000	14	
13	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Leaf curl (<i>Taphrina deformans</i> ; TAPHDE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	Not acceptable (worker, PPE*)
25	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Monilia (<i>Monilia</i> spp.; 1MONLG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
15	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Shot-hole (<i>Stigmia carpophila</i> ; STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
16	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
17	FR (SEZ)	Apricot (<i>Prunus armeniaca</i> ; 0140010; FS 0240)	F	Cytospora canker (<i>Cytospora leucostoma</i> ; VALSLE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
19	FR (SEZ)	Cherry (<i>Prunus avium</i> ; 0140020; FS 0013, FS0243, FS 0244)	F	Monilia (<i>Monilia</i> spp.; 1MONLG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	Not acceptable (worker, PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
20	FR (SEZ)	Cherry (<i>Prunus avium</i> ; 0140020; FS 0013, FS0243, FS 0244)	F	Shot-hole (<i>Stigmina carpophila</i> ; STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
21	FR (SEZ)	Cherry (<i>Prunus avium</i> ; 0140020; FS 0013, FS0243, FS 0244)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
23	FR (SEZ)	Cherry (<i>Prunus avium</i> ; 0140020; FS 0013, FS0243, FS 0244)	F	Anthrachnose (<i>Blumeriella jaapii</i> ; BLUMJA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
24	FR (SEZ)	Peach (<i>Prunus persica</i> ; 0140030; FS 2001, FS 0247) Nectarine (<i>Prunus persica</i> var. <i>nucipersica</i> ; 0140030; FS 2001, FS 0245)	F	Leaf curl (<i>Taphrina deformans</i> ; TAPHDE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	Not acceptable (worker, PPE*)
25	FR (SEZ)	Peach (<i>Prunus persica</i> ; 0140030; FS 2001, FS 0247) Nectarine (<i>Prunus persica</i> var. <i>nucipersica</i> ; 0140030; FS 2001, FS 0245)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
26	FR (SEZ)	Peach <i>(Prunus persica;</i> 0140030; FS 2001, FS 0247)	F	Shot-hole <i>(Stigmia carpophila;</i> STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
		Nectarine <i>(Prunus persica var.</i> <i>nucipersica;</i> 0140030; FS 2001, FS 0245)											
27	FR (SEZ)	Peach <i>(Prunus persica;</i> 0140030; FS 2001, FS 0247)	F	Bacterial canker <i>(Pseudomonas syringae</i> <i>pv. syringae;</i> PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
		Nectarine <i>(Prunus persica var.</i> <i>nucipersica;</i> 0140030; FS 2001, FS 0245)											
28	FR (SEZ)	Peach <i>(Prunus persica;</i> 0140030; FS 2001, FS 0247)	F	Cytospora canker <i>(Cytospora leucostoma;</i> VALSLE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
		Nectarine <i>(Prunus persica var.</i> <i>nucipersica;</i> 0140030; FS 2001, FS 0245)											
29	FR (SEZ)	Plum <i>(Prunus sp;</i> 0140040; FS 0014)	F	Leaf curl <i>(Taphrina deformans;</i> TAPHDE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	Not acceptable (worker, PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
30	FR (SEZ)	Plum (<i>Prunus sp</i> ; 0140040; FS 0014)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
31	FR (SEZ)	Plum (<i>Prunus sp</i> ; 0140040; FS 0014)	F	Shot-hole (<i>Stigmia carpophila</i> ; STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
32	FR (SEZ)	Plum (<i>Prunus sp</i> ; 0140040; FS 0014)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	F	
34	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	Scab (<i>Venturia inaequalis</i> ; VENTIN)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker, PPE*)
35	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	European canker (<i>Nectria galligena</i> ; NECTGA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
36	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
37	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	Blister spot (<i>Pseudomonas syringae</i> <i>pv. papulans</i> ; PSDMPA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
38	FR (SEZ)	Apple (<i>Malus domestica</i> ; 0130010; FP 0226)	F	Fire blight (<i>Erwinia amylovora</i> ; ERWIAM)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
39	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	Scab (<i>Venturia pyrina</i> ; VENTPI)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker, PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
40	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	European canker (<i>Nectria galligena</i> ; NECTGA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker, PPE*)
41	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
42	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
43	FR (SEZ)	Pear (<i>Pyrus communis</i> ; 0130020; FP 0230)	F	Fire blight (<i>Erwinia amylovora</i> ; ERWIAM)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
44	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	Scab (<i>Venturia pyrina</i> ; VENTPI)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
45	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	European canker (<i>Nectria galligena</i> ; NECTGA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
46	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
47	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
48	FR (SEZ)	Quince (<i>Cydonia oblonga</i> ; 0130030; FP 0231)	F	Fire blight (<i>Erwinia amylovora</i> ; ERWIAM)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
49	FR (SEZ)	Medlar and Japanese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	Scab (<i>Venturia inaequalis</i> ; VENTIN)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker, PPE*)
50	FR (SEZ)	Medlar and Japanese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	European canker (<i>Nectria galligena</i> ; NECTGA)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
51	FR (SEZ)	Medlar and Japanese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	Monilia (<i>Monilia spp.</i> ; 1MONLG)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
52	FR (SEZ)	Medlar and Japanese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
53	FR (SEZ)	Medlar and Japanese medlar (<i>Mespilus germanica</i> ; 0130040; FP 0229, <i>Eriobotrya japonica</i> , 0130050; FP 4044)	F	Fire blight (<i>Erwinia amylovora</i> ; ERWIAM)	Foliar spray	91-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
54	FR (SEZ)	Almond (<i>Prunus dulcis</i> ; 0120010; TN 0660)	F	Leaf curl (<i>Taphrina deformans</i> ; TAPHDE)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker, PPE*)
55	FR (SEZ)	Almond (<i>Prunus dulcis</i> ; 0120010; TN 0660)	F	Monilia (<i>Monilia spp.</i> ; 1MONIG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
56	FR (SEZ)	Almond (<i>Prunus dulcis</i> ; 0120010; TN 0660)	F	Shot-hole (<i>Stigmia carpophila</i> ; STIGCA)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
57	FR (SEZ)	Almond (<i>Prunus dulcis</i> ; 0120010; TN 0660)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
59	FR (SEZ)	Walnut (<i>Juglans regia</i> ; 0120110; TN 0678)	F	Alternaria (<i>Alternaria spp.</i> ; 1ALTEG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker, PPE*)
61	FR (SEZ)	Walnut (<i>Juglans regia</i> ; 0120110; TN 0678)	F	Bacterial blight (<i>Xanthomonas arboricola</i> <i>pv. juglandis</i> ; XANTJU)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
62	FR (SEZ)	Walnut (<i>Juglans regia</i> ; 0120110; TN 0678)	F	Pseudomonas (<i>Pseudomonas spp.</i> ; 1PSDMG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
64	FR (SEZ)	Hazelnut (<i>Corylus avellana</i> ; 0120060; TN 0666)	F	Alternaria (<i>Alternaria spp.</i> ; 1ALTEG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker, PPE*)
65	FR (SEZ)	Hazelnut (<i>Corylus avellana</i> ; 0120060; TN 0666)	F	Cytospora canker (<i>Cytospora spp.</i> ; 1CYTOG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
66	FR (SEZ)	Hazelnut (<i>Corylus avellana</i> ; 0120060; TN 0666)	F	Bacterial blight (<i>Xanthomonas arboricola</i> <i>pv. corylina</i> ; XANTCY)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
67	FR (SEZ)	Hazelnut (<i>Corylus avellana</i> ; 0120060; TN 0666)	F	Stem dieback (<i>Pseudomonas avellanae</i> ; PSDMAL)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
68	FR (SEZ)	Chestnut (<i>Castanea sativa</i> ; 0120040; TN 0664)	F	Mycosphaerella (<i>Mycosphaerella spp.</i> ; 1MYCOG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker, PPE*)
69	FR (SEZ)	Chestnut (<i>Castanea sativa</i> ; 0120040; TN 0664)	F	Alternaria (<i>Alternaria spp.</i> ; 1ALTEG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
71	FR (SEZ)	Chestnut (<i>Castanea sativa</i> ; 0120040; TN 0664)	F	Xanthomonas (<i>Xanthomonas spp.</i> ; 1XANTG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
72	FR (SEZ)	Chestnut (<i>Castanea sativa</i> ; 0120040; TN 0664)	F	Pseudomonas (<i>Pseudomonas spp.</i> ; 1PSDMG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
74	FR (SEZ)	Pistachio (<i>Pistacia vera</i> ; 0120100; TN 0675)	F	Alternaria (<i>Alternaria spp.</i> ; 1ALTEG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	Not acceptable (worker, PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
76	FR (SEZ)	Pistachio (<i>Pistacia vera</i> ; 0120100; TN 0675)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
77	FR (SEZ)	Pistachio (<i>Pistacia vera</i> ; 0120100; TN 0675)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	95-53	a) 4 b) 4	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	500- 1000	F	
78	FR (SEZ)	Olive (Table olives, olive for oil production) (<i>Olea europaea</i> ; 0161030, 0402010; FT 0305, SO 0305)	F	Peacock spot (<i>Spilocaea oleaginea</i> ; CYCLOL)	Foliar spray	10-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	600- 1000	14	Not acceptable (worker, PPE*) <i>Efficacy demonstrated on anthracnose</i>
79	FR (SEZ)	Olive (Table olives, olive for oil production) (<i>Olea europaea</i> ; 0161030, 0402010; FT 0305, SO 0305)	F	Bacterial canker (<i>Pseudomonas savastoniai</i> <i>pv. savastoniai</i> ; PSDMSA)	Foliar spray	10-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	600- 1000	14	
80	FR (SEZ)	Olive (Table olives, olive for oil production) (<i>Olea europaea</i> ; 0161030, 0402010; FT 0305, SO 0305)	F	Anthracnose (<i>Colletotrichum</i> <i>gloeosporioides</i> ; GLOMCI)	Foliar spray	10-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	600- 1000	14	
81	FR (SEZ)	Grape (Table grape, wine grapes) (<i>Vitis vinifera</i> ; 0151000, 0151010, 0151020; FB 0269)	F	Downy mildew (<i>Plasmopara viticola</i> ; PLASVI)	Foliar spray	15-81 & 91	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	21	Not acceptable (worker, PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
83	FR (SEZ)	Grape (Table grape, wine grapes) <i>(Vitis vinifera</i> ; 0151000, 0151010, 0151020; FB 0269)	F	Bacterial blight (<i>Xanthomonas ampelina</i> ; XANTAM)	Foliar spray	15-81 & 91	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	21	
84	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Late blight (<i>Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	Not acceptable (open- field and walk-in tunnels) (PPE*)
85	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
86	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Anthraxnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
87	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
88	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	F	Bacterial speck (<i>Pseudomonas syringae pv. tomato</i> ; PSDMTM)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
89	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	F	Late blight (<i>Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	Not acceptable (open- field and walk-in tunnels) (PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
90	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	F	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
91	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	F	Anthrachnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
92	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	F	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
93	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	F	Bacterial speck (<i>Pseudomonas syringae pv. tomato</i> ; PSDMTM)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	10	
94	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Blight of pepper (<i>Phytophthora capsici</i> ; PHYTCP)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL) Not acceptable (open- field and walk-in tunnel) (PPE*)
95	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
96	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Anthrachnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
97	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
98	FR (SEZ)	Pepper (<i>Capsicum annuum</i> ; 0231020; VO 0051)	F	Bacterial canker (<i>Pseudomonas syringae</i> <i>pv. syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
99	FR (SEZ)	Leek (<i>Allium porrium</i> ; 0270060; VA 0384) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Downy mildew (<i>Phytophthora porri</i> ; PHYTPO)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL, PPE*)
104	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Downy mildew (<i>Bremia lactucae</i> ; BREMLA)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL, PPE*)
105	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Ascochyta (<i>Ascochyta spp</i> ; 1ASCOG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
106	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
107	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
108	FR (SEZ)	Artichoke (<i>Cynara scolymus</i> ; 0270050; VS 0620) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
109	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Downy mildew (<i>Plasmopara crustosa</i> ; PLASCR)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL, PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
110	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Ascochyta (<i>Ascochyta spp</i> ; (1ASCOG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
111	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
112	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
113	FR (SEZ)	Celery (<i>Apium graveolens</i> ; 0270030; VS 0624) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
114	FR (SEZ)	Cardoon (<i>Cynara cardunculus</i> ; 0270020; VS 0623) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Downy mildew (<i>Bremia lactucae</i> ; BREMLA)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL, PPE*)
115	FR (SEZ)	Cardoon (<i>Cynara cardunculus</i> ; 0270020; VS 0623) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Ascochyta (<i>Ascochyta spp</i> ; 1ASCOG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
116	FR (SEZ)	Cardoon (<i>Cynara cardunculus</i> ; 0270020; VS 0623) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
117	FR (SEZ)	Cardoon (<i>Cynara cardunculus</i> ; 0270020; VS 0623) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
118	FR (SEZ)	Cardoon (<i>Cynara cardunculus</i> ; 0270020; VS 0623) Whole group of stem vegetables except asparagus (0270000; VS 0078)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-51	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
119	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	Downy mildew (<i>Hyaloperonospora brassicae</i> ; HPERBR)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL, PPE*)
120	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	Black spot (<i>Alternaria brassicae</i> ; ALTEBA)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
121	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	<i>Colletotrichum</i> (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
122	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	Black rot (<i>Xanthomonas campestris pv. campestris</i> ; XANTCA)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
123	FR (SEZ)	Broccoli (<i>Brassica oleracea</i> ; 0241010; VB 0400, VB 0401)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
124	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Downy mildew (<i>Hyaloperonospora</i> <i>brassicae</i> ; HPERBR)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL, PPE*)
125	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Black spot (<i>Alternaria brassicae</i> ; ALTEBA)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
126	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Colletotrichum (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
127	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Black rot (<i>Xanthomonas campestris</i> pv. <i>campestris</i> ; XANTCA)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
128	FR (SEZ)	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i> ; 0241020; VB 0404)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-59	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
129	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (open- field and walk-in tunnel) (PPE*)
130	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
131	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
132	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
133	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
134	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (open- field and walk-in tunnel) (PPE*)
135	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
136	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Anthrachnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
137	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
138	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
139	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not finalised (open- field and walk-in tunnel) (analytical methods, consumer exposure, birds, mammals, aquatic organisms, bees and bumble bees, soil macro-organisms, PPE*)
140	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
141	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Anthraco nose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
142	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
143	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
144	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL)
145	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (open- field and walk-in tunnel) (PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
146	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLIA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
147	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
148	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
149	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL) Not acceptable (open- field and walk-in tunnel) (PPE*)
150	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
151	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLIA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
152	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
153	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	F	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
154	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
155	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (open- field and walk-in tunnel) (PPE*)
156	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Anthraco nose (<i>Colletotrichum lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
157	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Bacterial leaf spot (<i>Xanthomonas campestris pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
158	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	F	Angular leaf spot (<i>Pseudomonas syringae pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
159	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	Downy mildew (<i>Peronospora destructor</i> ; PERODE)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
160	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	Purple blotch (<i>Alternaria porri</i> ; ALTEPO)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
161	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	Anthraco nose (<i>Colletotrichum coccodes</i> ; COLLCC)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
162	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	Leaf blight (<i>Pleospora allii</i> ; PLEOAL)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
163	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
164	FR (SEZ)	Onion (<i>Allium cepa</i> ; 0220020; VA 0385, VA 0386, VA 0387)	F	<i>Xanthomonas</i> (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
165	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	Downy mildew (<i>Peronospora destructor</i> ; PERODE)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
166	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	Purple blotch (<i>Alternaria porri</i> ; ALTEPO)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
167	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	Antracnose (<i>Colletotrichum coccodes</i> ; COLLCC)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
168	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	Leaf blight (<i>Pleospora allii</i> ; PLEOAL)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
169	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
170	FR (SEZ)	Garlic (<i>Allium sativum</i> ; 0220010; VA 0381)	F	<i>Xanthomonas</i> (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
171	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Downy mildew (<i>Peronospora destructor</i> ; PERODE)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
172	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Purple blotch (<i>Alternaria porri</i> ; ALTEPO)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
173	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Antracnose (<i>Colletotrichum coccodes</i> ; COLLCC)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
174	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Leaf blight (<i>Pleospora allii</i> ; PLEOAL)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
175	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
176	FR (SEZ)	Shallot (<i>Allium cepa</i> <i>Aggregatum</i> types; 0220030; VA 0388)	F	Xanthomonas (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	14-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
177	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	F	Downy mildew (<i>Bremia lactucae</i> ; BREMLA)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL) Not acceptable (open- field and walk-in tunnel) (PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
178	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	F	Leaf spot (<i>Alternaria sonchi</i> ; ALTESN)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable against grey mould and sclerotinia (efficacy)
179	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	F	Anthrachnose (<i>Marssonina panattoniana</i> ; MARSPA)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
180	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	F	Sclerotinia (<i>Sclerotinia minor</i> ; SCLEMI)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
181	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
182	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	F	Bacterial spot (<i>Xanthomonas axonopodis</i> pv. <i>vitians</i> ; XANTVI)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
183	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and similar (leaves) (0252000)	F	Downy mildew (<i>Peronospora farinosa f.</i> <i>sp. Spinaciae</i> ;PEROFS)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL)
184	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and similar (leaves) (0252000)	F	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (open- field and walk-in tunnel) (PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
185	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and similar (leaves) (0252000)	F	Anthrachnose (<i>Colletotrichum spinaciae</i> ; COLLDS)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
188	FR (SEZ)	Bean (Bean with pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Downy mildew (<i>Phytophthora phaseoli</i> ; PHYTPH)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*) Not acceptable against rust (efficacy)
189	FR (SEZ)	Bean (Bean with pods, bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Anthrachnose (<i>Colletotrichum lindemuthianum</i> ; COLLLD)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
190	FR (SEZ)	Bean (Bean with pods, bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Rust (<i>Uromyces appendiculatus</i> ; UROMAP)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
191	FR (SEZ)	Bean (Bean with pods, bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Septoria (<i>Septoria spp</i> ; 1SEPTG)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
192	FR (SEZ)	Bean (Bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	Bacterial blight (<i>Xanthomonas axonopodis</i> <i>pv. phaseoli</i> ; XANTPH)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
193	FR (SEZ)	Bean (Bean without pods) (<i>Phaseolus vulgaris</i> ; 0260010. 0260020; VP 0061, <i>Vicia faba</i> ; 0260020; VP 4407, VP0522, VP 0523)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
194	FR (SEZ)	Pea (Pea with pods) (<i>Pisum sativum</i> ; 0260030, 0260040; VP 4441)	F	Downy mildew (<i>Peronospora viciae</i> ; PEROVI)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable for peas without pods (MRL) Not acceptable (PPE*) Not acceptable against rust (efficacy)
195	FR (SEZ)	Pea (Pea with pods, pea without pods) (<i>Pisum sativum</i> ; 0260030, 0260040; VP 4441)	F	Anthrachnose (<i>Colletotrichum</i> <i>lindemuthianum</i> ; COLLLD)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
196	FR (SEZ)	Pea (Pea with pods, pea without pods) (<i>Pisum sativum</i> ; 0260030, 0260040; VP 4441)	F	Rust (<i>Uromyces fabae</i> ; UROMVF)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (MRL) Not acceptable (PPE*) Not acceptable against rust (efficacy)
197	FR (SEZ)	Pea (Pea with pods, pea without pods) (<i>Pisum sativum</i> ; 0260030, 0260040; VP 4441)	F	Leaf blotch (<i>Septoria pisi</i> ; SEPTPI)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
201	FR (SEZ)	Lentils (<i>Lens culinaris</i> ; 0260050; VP 0533)	F	Anthrachnose (<i>Colletotrichum</i> <i>lindemuthianum</i> ; COLLLD)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
202	FR (SEZ)	Lentils (<i>Lens culinaris</i> ; 0260050; VP 0533)	F	Rust (<i>Uromyces fabae</i> ; UROMVF)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
203	FR (SEZ)	Lentils (<i>Lens culinaris</i> ; 0260050; VP 0533)	F	Septoria (<i>Septoria spp</i> ; 1SEPTG)	Foliar spray	14-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
206	FR (SEZ)	Potato (<i>Solanum</i> <i>tuberosum</i> ; 0211000; VR 0589)	F	Late blight (<i>Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL, PPE*)
207	FR (SEZ)	Potato (<i>Solanum</i> <i>tuberosum</i> ; 0211000; VR 0589)	F	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
208	FR (SEZ)	Potato (<i>Solanum tuberosum</i> ; 0211000; VR 0589)	F	Anthrachnose (<i>Colletotrichum coccodes</i> ; COLLCC)	Foliar spray	15-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
211	FR (SEZ)	Carrot (<i>Daucus carota subsp. Sativus</i> ; 0213020; VR 0577) Other root and tuber vegetables except sugar beet (0213000)	F	Downy mildew (<i>Plasmopara nivea</i> ; PLASCR)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable for carrot only (MRL) Not acceptable (PPE*)
212	FR (SEZ)	Carrot (<i>Daucus carota subsp. Sativus</i> ; 0213020; VR 0577) Other root and tuber vegetables except sugar beet (0213000)	F	Leaf blight (<i>Alternaria dauci</i> ; ALTEDA)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
213	FR (SEZ)	Carrot (<i>Daucus carota subsp. Sativus</i> ; 0213020; VR 0577) Other root and tuber vegetables except sugar beet (0213000)	F	Anthrachnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
214	FR (SEZ)	Carrot (<i>Daucus carota</i> <i>subsp. Sativus</i> ; 0213020; VR 0577) Other root and tuber vegetables except sugar beet (0213000)	F	Brown leaf spot (<i>Cercospora carotae</i> ; CERCCA)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
215	FR (SEZ)	Carrot (<i>Daucus carota</i> <i>subsp. Sativus</i> ; 0213020; VR 0577) Other root and tuber vegetables except sugar beet (0213000)	F	Bacterial leaf blight (<i>Xanthomonas campestris</i> <i>pv. carotae</i> ; XANTCR)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
216	FR (SEZ)	Carrot (<i>Daucus carota</i> <i>subsp. Sativus</i> ; 0213020; VR 0577) Other root and tuber vegetables except sugar beet (0213000)	F	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	15-47	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
217	FR (SEZ)	Sugar beet (<i>Beta vulgaris subsp.</i> <i>vulgaris</i> ; 0900010; VR 0596)	F	Leaf spot (<i>Cercospora beticola</i> ; CERCBE)	Foliar spray	39-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL) Not finalised (efficacy,, PPE*)
218	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Anthrachnose (<i>Colletotrichum fragariae</i> ; COLLFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
219	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Leaf spot (<i>Mycosphaerella fragariae</i> ; MYCOFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (open- field and walk-in tunnel) (PPE*)
220	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Red stele (<i>Phytophthora fragariae</i> ; PHYTFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
221	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Angular leaf spot (<i>Xanthomonas fragariae</i> ; XANTFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
222	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	F	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
223	FR (SEZ)	Ornamental plants and trees	F	Peronospora (<i>Peronospora spp</i> ; 1PEROG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	Not acceptable (open- field and walk-in tunnel) (worker, PPE*)
224	FR (SEZ)	Ornamental plants and trees	F	Septoria (<i>Septoria spp</i> ; 1SEPTG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
225	FR (SEZ)	Ornamental plants and trees	F	Colletotrichum (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	Not acceptable against rust (efficacy)
226	FR (SEZ)	Ornamental plants and trees	F	Puccinia (<i>Puccinia spp</i> ; 1PUCCG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
227	FR (SEZ)	Ornamental plants and trees	F	Monilia (<i>Monilia spp</i> ; 1MONLG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
Interzonal uses (use as seed treatment, in greenhouses (or other closed places of plant production), as post-harvest treatment or for treatment of empty storage rooms)													
228	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Late blight (<i>Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
229	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
230	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Anthracnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
231	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
232	FR (SEZ)	Tomato (<i>Solanum lycopersicum</i> ; 0231010; VO 0448)	G	Bacterial speck (<i>Pseudomonas syringae pv. tomato</i> ; PSDMTM)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
233	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Late blight (<i>Phytophthora infestans</i> ; PHYTIN)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
234	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
235	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Anthracnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
236	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
237	FR (SEZ)	Aubergine (Eggplant) (<i>Solanum melongena</i> ; 0231030; VO 0440)	G	Bacterial speck (<i>Pseudomonas syringae pv. tomato</i> ; PSDMTM)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
238	FR (SEZ)	Pepper (<i>Capsicum annum</i> ; 0231020; VO 0051)	G	Blight of pepper (<i>Phytophthora capsici</i> ; PHYTCP)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	Not acceptable (MRL,PPE*)
239	FR (SEZ)	Pepper (<i>Capsicum annum</i> ; 0231020; VO 0051)	G	Early blight (<i>Alternaria solani</i> ; ALTESO)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
240	FR (SEZ)	Pepper (<i>Capsicum annum</i> ; 0231020; VO 0051)	G	Anthracnose (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
241	FR (SEZ)	Pepper (<i>Capsicum annum</i> ; 0231020; VO 0051)	G	Bacterial spot (<i>Xanthomonas spp</i> ; 1XANTG)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
242	FR (SEZ)	Pepper (<i>Capsicum annum</i> ; 0231020; VO 0051)	G	Bacterial canker (<i>Pseudomonas syringae pv. syringae</i> ; PSDMSY)	Foliar spray	15-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	14	
243	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Downy mildew (<i>Pseudoperonospora cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha (f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
244	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
245	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Anthraxnose (<i>Colletotrichum lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
246	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Bacterial leaf spot (<i>Xanthomonas campestris pv. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
247	FR (SEZ)	Cucumber (<i>Cucumis sativus</i> ; 0232010; VC 0424)	G	Angular leaf spot (<i>Pseudomonas syringae pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
248	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Downy mildew (<i>Pseudoperonospora cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
249	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
250	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
251	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
252	FR (SEZ)	Courgette (zucchini) (<i>Cucurbita pepo</i> var. <i>giromontiina</i> ; 0232030; VC 4207, VC 4263)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
253	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (PPE*)
254	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
255	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	
256	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
257	FR (SEZ)	Gherkin (<i>Cucumis sativus</i> ; 0232020; VC 0425)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	3	Not acceptable (MRL, PPE*)
258	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
259	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
260	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Anthraco nose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
261	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
262	FR (SEZ)	Melon (<i>Cucumis melo</i> ; 0233010; VC 0046)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
263	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL, PPE*)
264	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
265	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLIA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
266	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
267	FR (SEZ)	Watermelon (<i>Citrullus lanatus</i> ; 0233030; VC 0432)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
268	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Downy mildew (<i>Pseudoperonospora</i> <i>cubensis</i> ; PSPECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL, PPE*)
269	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Leaf blight (<i>Alternaria cucumerina</i> ; ALTECU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
270	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Anthracnose (<i>Colletotrichum</i> <i>lagenarium</i> ; COLLIA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
271	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Bacterial leaf spot (<i>Xanthomonas campestris</i> <i>pc. cucurbitae</i> ; XANTCU)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
272	FR (SEZ)	Pumpkin (marrow) (<i>Cucurbita maxima</i> ; 0233020; VC 0429)	G	Angular leaf spot (<i>Pseudomonas syringae</i> <i>pv. lachrymans</i> ; PSDMLA)	Foliar spray	10-89	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
273	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	G	Downy mildew (<i>Bremia lactucae</i> ; BREMLA)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (PPE*) Not acceptable against grey mould and sclerotinia (efficacy)
274	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	G	Leaf spot (<i>Alternaria sonchi</i> ; ALTESN)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
275	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	G	Anthrachnose (<i>Marssonina panattoniana</i> ; MARSPA)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
276	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	G	Sclerotinia (<i>Sclerotinia minor</i> ; SCLEMI)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
277	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	G	Pseudomonas (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
278	FR (SEZ)	Lettuce (<i>Lactuca sativa</i> ; 0251020; VL 0510, VL 0482, VL 0483) Lettuce and other salad plants including leaves and sprouts of brassicacea (0251000)	G	Bacterial spot (<i>Xanthomonas axonopodis</i> <i>pv. vitians</i> ; XANTVI)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/ma x		
279	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and similar (leaves) (0252000)	G	Downy mildew (<i>Peronospora farinosa</i> f. <i>sp. Spinaciae</i> ;PEROFS)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL, PPE*)
280	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and similar (leaves) (0252000)	G	Alternaria (<i>Alternaria spp</i> ; 1ALTEG)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
281	FR (SEZ)	Spinach (<i>Spinacia oleracea</i> ; 0252010; VL 0502) Spinach and similar (leaves) (0252000)	G	Anthrachnose (<i>Colletotrichum spinaciae</i> ; COLLDS)	Foliar spray	12-49	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
284	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	Anthrachnose (<i>Colletotrichum fragariae</i> ; COLLFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	Not acceptable (MRL, PPE*)
285	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	Leaf spot (<i>Mycosphaerella fragariae</i> ; MYCOFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
286	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	Red stele (<i>Phytophthora fragariae</i> ; PHYTFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
287	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	Angular leaf spot (<i>Xanthomonas fragariae</i> ; XANTFR)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. ^(e)	Member state(s)	Crop and/ or situation (crop destination/purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks: e.g. g safener/synergist per ha ^(f)
					Method/Ki nd	Timing/Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	kg product/ha a) max. rate per appl. b) max. total rate per crop/season	g a.s./ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha min/max		
288	FR (SEZ)	Strawberry (<i>Fragaria spp</i> ; 0152000; FB0275)	G	<i>Pseudomonas</i> (<i>Pseudomonas spp</i> ; 1PSDMG)	Foliar spray	13-85	a) 5 b) 5	7 7	a) 5.0 b) 20.0	a) 1.0 b) 4.0	400- 1000	7	
289	FR (SEZ)	Ornamental plants and trees	G	<i>Peronospora</i> (<i>Peronospora spp</i> ; 1PEROG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	Not acceptable (PPE*) Not acceptable against rust (efficacy)
290	FR (SEZ)	Ornamental plants and trees	G	<i>Septoria</i> (<i>Septoria spp</i> ; 1SEPTG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
291	FR (SEZ)	Ornamental plants and trees	G	<i>Colletotrichum</i> (<i>Colletotrichum spp</i> ; 1COLLG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
292	FR (SEZ)	Ornamental plants and trees	G	<i>Puccinia</i> (<i>Puccinia spp</i> ; 1PUCCG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	
293	FR (SEZ)	Ornamental plants and trees	G	<i>Monilia</i> (<i>Monilia spp</i> ; 1MONLG)	Foliar spray	All stages	a) 3 b) 3	7 7	a) 5.0 b) 15.0	a) 1.0 b) 3.0	400- 1000	NA	

* PPE : Personal protective equipment. The information is insufficient to ensure the performance of the clothing PPE (in accordance with standard NF EN ISO 27065/A1) for this type of formulation (WP). A. Shaw *and al.* (2018).

Remarks table heading:

(a) e.g. wettable powder (WP), emulsifiable concentrate (EC), granule (GR)
(b) Catalogue of pesticide formulation types and international coding system CropLife International Technical Monograph n°2, 6th Edition Revised May 2008
(c) g/kg or g/l

(d) Select relevant
(e) Use number(s) in accordance with the list of all intended GAPs in Part B, Section 0 should be given in column 1
(f) No authorisation possible for uses where the line is highlighted in grey, Use should be crossed out when the notifier no longer supports this use.

Super Bouillie Macclesfield 80

Part A - National Assessment

FRANCE DEPR version

Remarks	1	Numeration necessary to allow references	7	Growth stage at first and last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application
columns:	2	Use official codes/nomenclatures of EU Member States	8	The maximum number of application possible under practical conditions of use must be provided.
	3	For crops, the EU and Codex classifications (both) should be used; when relevant, the use situation should be described (e.g. fumigation of a structure)	9	Minimum interval (in days) between applications of the same product
	4	F: professional field use, Fn: non-professional field use, Fpn: professional and non-professional field use, G: professional greenhouse use, Gn: non-professional greenhouse use, Gpn: professional and non-professional greenhouse use, I: indoor application	10	For specific uses other specifications might be possible, e.g.: g/m ³ in case of fumigation of empty rooms. See also EPPO-Guideline PP 1/239 Dose expression for plant protection products.
	5	Scientific names and EPPO-Codes of target pests/diseases/ weeds or, when relevant, the common names of the pest groups (e.g. biting and sucking insects, soil born insects, foliar fungi, weeds) and the developmental stages of the pests and pest groups at the moment of application must be named.	11	The dimension (g, kg) must be clearly specified. (Maximum) dose of a.s. per treatment (usually g, kg or L product/ha).
	6	Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plants - type of equipment used must be indicated.	12	If water volume range depends on application equipments (e.g. ULVA or LVA) it should be mentioned under "application: method/kind".
			13	PHI - minimum pre-harvest interval
			14	Remarks may include: Extent of use/economic importance/restrictions

3 Background of authorisation decision and risk management

3.1 Physical and chemical properties (Part B, Section 2)

SUPER SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) is a solid wettable powder (WP). All studies have been performed in accordance with the current requirements and the results are deemed acceptable. The appearance of the product is a medium turquoise solid powder. The product is not explosive and has no oxidising properties. It is not flammable. It has no self-ignition temperature below 400°C. In aqueous solution 1%, it has a pH value of 6.75 at 22°C. There is no effect of high temperature on the stability of the formulation, since after 14 days at 54 °C, neither the active ingredient content nor the technical properties were changed. The stability data indicate a shelf life of at least two years at ambient temperature when stored in LDPE and HDPE, PE. Its technical characteristics are acceptable for a WP formulation.

The formulation is not classified for the physico-chemical aspect.

When water soluble bags are used, the slurry should be stirred for at least 3 minutes before application.

3.2 Efficacy (Part B, Section 3)

Considering the data submitted:

- The efficacy level of SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) is considered acceptable for most of the requested uses. **For uses intended to control grey mould and sclerotinia on lettuce, rusts on beans, peas, floral crops and green plants, and cercosporiosis on garden beets, the demonstration of the efficacy of the product is considered insufficient.**
- The phytotoxicity level of SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) is considered acceptable for all the requested uses. On ornamental crops, considering the diversity of crops, it is recommended, in case of doubts, to test the product selectivity on a small area. Moreover, it is also recommended to avoid applications during flowering, because of a risk of burnings on flowers.
- The risks of negative impact on yield, cider making process, propagation, succeeding crops, adjacent crops are considered negligible. **Risks with copper such as spotting of table grape berries or on the wine-making process are known.** However, these risks of negative impact are considered acceptable.
- There is a risk of resistance developing or appearing to copper for *Xanthomonas* bacteria that requires a monitoring on vegetable crops (e.g. tomato) and walnut.

3.3 Methods of analysis (Part B, Section 5)

3.3.1 Analytical method for the formulation

Analytical methods for the determination of the active substance and the relevant impurities in the formulation are available and validated.

3.3.2 Analytical methods for residues

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Analytical methods are available in the Draft Assessment Report and in this dossier, and are validated for the determination of residues of copper in plants (high water, oily, acidic and dry content commodities), soil, water (surface and drinking), air and body fluids.

According to EFSA conclusions, an ILV of the analytical methods for the determination of residues of copper in plants is required.

Analytical methods for the determination of residues of copper in foodstuffs of animal origin are missing and are required. Moreover, the LOQ of the available methods for the determination of residues of copper in water is not in accordance with the European Directive 98/83/EC.

3.4 Mammalian toxicology (Part B, Section 6)

Endpoints used in risk assessment

Agreed EU endpoints	
Active substance	Copper (Bordeaux mixture) 200 g/kg
AOEL systemic	0.08 mg/kg bw/d
AAOEL	Not necessary
Inhalation absorption	100%
Oral absorption	50%
Vapour pressure	Not necessary
Dermal absorption	Concentrate: 1% Dilution: 9% (Based on <i>in vitro</i> through human skin studies)

3.4.1 Acute toxicity

SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80), containing 200 g/kg copper (under the form of Bordeaux mixture), has a low acute oral and dermal toxicity, is harmful if inhaled, is not irritating to the rabbit skin or eye and is not a skin sensitiser.

3.4.2 Operator exposure

Considering the proposed uses, operator systemic exposure was estimated using the EFSA model¹¹:

Outdoor:

		Copper under the form of Bordeaux mixture	
Model data	Level of PPE	Total absorbed dose (mg/kg/day)	% of systemic AOEL
Critical use: Olive (worst-case scenario for high crops) ¹			

¹¹ AOEM – Agricultural Operator Exposure Model (EFSA Journal 2014;12 (10):3874)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Tractor mounted outdoor, upward applications			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile) Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and A	0.0976	122
	Work wear (arms, body and legs covered) and gloves M/L and A and mask (FP1, P1 and similar) during M/L	0.0437	55
	Work wear (arms, body and legs covered) and gloves M/L and A and closed cab during A	0.0764	96
	Work wear (arms, body and legs covered) and gloves M/L and A; mask (FP1, P1 and similar) during M/L and closed cab during A	0.0225	28
Manual knapsack outdoor, upward applications (late: dense foliage of the crop) ³			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile) Body weight: 60 kg	Work wear (arms, body and legs covered) M/L and A	0.1020	127.5
	Work wear (arms, body and legs covered) and gloves M/L and A	0.0734	92
Critical use: Ornamentals plants and trees			
Manual Handheld, outdoor, upward applications (late: dense foliage of the crop)			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile) Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves during M/L and A	0.1095	137
	Work wear (arms, body and legs covered) and gloves M/L and A and mask (FP1, P1 and similar) during M/L	0.0823	103
	Work wear (arms, body and legs covered), gloves and mask (FP1, P1 and similar) during M/L and A	0.0799	99.9
	Work wear (arms, body and legs covered) and gloves M/L and A and mask (FP2, P2 and similar) during M/L	0.0769	96
Critical use: Broccoli (worst-case scenario for low crops) ²			
Tractor mounted outdoor, downward applications			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile) Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and A	0.1257	157
	Work wear (arms, body and legs covered) and gloves M/L and A and mask (FP1, P1 and similar) during M/L	0.0385	48

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Critical use: Ornamentals plants			
Manual hand held outdoor, downward applications			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile) Body weight: 60 kg	Work wear (arms, body and legs covered) and gloves M/L and A	0.0925	116
	Work wear (arms, body and legs covered) and gloves M/L and A and mask (FP1, P1 and similar) during M/L	0.0515	64
Manual knapsack, outdoor, downward applications			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile) Body weight: 60 kg	Work wear (arms, body and legs covered) M/L and A	0.0181	23

¹ covers grapes, pome fruits, stone fruits, tree nuts, olive tree, citrus trees and ornamental plants and trees (high crops).

² covers other flowering brassica, tomato, eggplant, pepper, fresh legumes, sugarbeet, stem vegetables, cucurbits edible peel, cucurbits inedible peel, bulb vegetables, lettuce, spinach, beans, potato, carrots, strawberry, and ornamental plants and trees (low crops).

Indoor:

Copper under the form of Bordeaux mixture			
Model data	Level of PPE	Total absorbed dose (mg/kg/day)	% of systemic AOEL
Critical use: Tomato ¹			
Manual Hand held, upward spraying			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile); 60 kg 1 ha/day	Potential exposure	0.2319	294
	Work wear (arms, body and legs covered) M/L/A	0.0862	108
	Work wear (arms, body and legs covered) and gloves M/L/A	0.0620	77.5
Manual knapsack, upward spraying			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile); 60 kg 1 ha/day	Potential exposure	0.1000	125
	Work wear (arms, body and legs covered) M/L/A	0.0092	11.5
Critical use: Lettuce ²			
Manual Hand held, downward spraying			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile); 60 kg 1 ha/day	Potential exposure	0.4659	582
	Work wear (arms, body and legs covered) M/L/A	0.1106	138
	Work wear (arms, body	0.0926	116

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

	and legs covered) and gloves M/L/A		
	Work wear (arms, body and legs covered) and gloves M/L/A, mask (FP1, P1 and similar) during M/L	0.0515	64
Manual knapsack, downward spraying			
Application rate		1 kg a.s./ha	
Spray application (AOEM; 75 th percentile); 60 kg 1 ha/day	Potential exposure	0.1382	173
	Work wear (arms, body and legs covered) M/L/A	0.0181	23

¹ covers eggplant, pepper, zucchini, cucumber, gherkin, strawberry and ornamentals

² covers melon, watermelon, pumpkin, spinach and ornamentals

Conclusion:

According to the EFSA model calculations, it can be concluded that the risk for the operator using SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) is **acceptable** taking into account the above mentioned personal protective equipment (PPE).

For details of personal protective equipment for workers, refer to the Decision in Appendix 1.

3.4.3 Worker exposure

Workers may have to enter in treated areas after treatment for crop inspection/irrigation, cutting, sorting, bundling, carrying, hand harvesting, reaching, picking and searching activities. Therefore, estimation of worker exposure was calculated according to the EFSA model.

Outdoor:

		Copper under the form of Bordeaux mixture	
Model data	Level of PPE	Total absorbed dose (mg/kg bw/day)	% of systemic AOEL
Critical Use: Grapes			
Hand harvesting			
Outdoor			
Work rate: 8 hours/day			
DT ₅₀ : 7 days			
DFR: 3µg/cm ² /kg a.s./ha			
Interval between applications: 7 days			
Application rate		4 x 1 kg.as/ha (5 x 0.8 kg.as/ha)	
EFSA model 2014 (AOEM Excel calculator) Body weight: 60 kg	Potential TC: 30000 cm ² /person/h	2.0250 (1.6740)	2531 (2093)
	Work wear (arms, body and legs covered) TC: 10100 cm ² /person/h	0.6818 (0.5636)	852 (704)
	Work wear (arms, body and legs covered) and gloves TC: No TC available for this	-	-

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

	assessment ¹²		
Critical Use: Citrus fruit Covering: olive, pome fruit, stone fruit and tree nuts			
Searching, reaching, picking Outdoor Work rate: 8 hours/day, DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days			
Number of applications and application rate		4 x 1 kg.as/ha (5 x 0.8 kg.as/ha ; 3 x 1 kg.as/ha)	
EFSA model 2014 (AOEM Excel calculator) Body weight: 60 kg	Potential TC: 22500 cm ² /person/h	1,5188 (1.2555; 1.4175)	1898 (1569; 1772)
	Work wear (arms, body and legs covered) TC: 4500 cm ² /person/h	0,3038 (0.2511; 0.2835)	380 (314; 354)
	Work wear (arms, body and legs covered) and gloves TC: 2250 cm ² /person/h	0,1519 (0.1256; 0.1418)	190 (157; 177)
Critical use: Cucumber Covering: tomato, eggplant, pepper, artichoke, zucchini, gherkin, melon, watermelon, pumpkin, leek, cardoon, lettuce, spinach, broccoli, cauliflower, onion, garlic, shallot, bean, pea, lentils.			
Reaching, picking Outdoor Work rate: 8 hours/day, DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days			
Number of applications and application rate		4 x 1 kg.as/ha (5 x 0.8 kg.as/ha as worst case comparing to 4 x 0.8 kg.as/ha or 3 x 0.8 kg.as/ha)	
EFSA model 2014 (AOEM Excel calculator) Body weight: 60 kg	Potential TC: 5800 cm ² /person/h	0,3915 (0.3236)	489 (405)
	Work wear (arms, body and legs covered) TC: 2500 cm ² /person/h	0,1688 (0.1395)	211 (174)
	Work wear (arms, body and legs covered) and gloves TC: 580 cm ² /person/h	0,0392 (0.0324)	49 (40.5)
Critical Use: Strawberry			
Reaching, picking Outdoor Work rate: 8 hours/day, DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days			
Number of applications and application rate		4 x 1 kg. as/ha	

¹² For grapes, the applicant proposed a specific refined TC value considering gloves. This TC value cannot be accepted since the setting of a new TC value at European level requires the analysis of all available data.

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

		(5 x 0.8 kg.a/ha as worst case comparing to 4 x 0.8kg.as/ha)	
EFSA model 2014 (AOEM Excel calculator) Body weight: 60 kg	Potential TC: 5800 cm ² /person/h	0,3915 (0.3236)	489 (404.5)
	Work wear (arms, body and legs covered) TC: 3000 cm ² /person/h	0,2025 (0.1674)	253 (209)
	Work wear (arms, body and legs covered) and gloves TC: 750 cm ² /person/h	0,0506 (0.0419)	63 (52)
Critical Use: Potato Covering: sugar beet, carrot, celery			
Inspection, irrigation Outdoor Work rate: 2 hours/day, DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days			
Number of applications and application rate		4 x 1 kg.as/ha (5 x 0.8 kg.as/ha as worst case comparing to 4 x 0.8kg.as/ha or 3 x 0.8 kg.as/ha)	
EFSA model 2014 (AOEM Excel calculator) Body weight: 60 kg	Potential TC: 12500 cm ² /person/h	0,2109 (0.1744)	264 (218)
	Work wear (arms, body and legs covered) TC: 1400 cm ² /person/h	0,0236 (0.0195)	30 (24)
	Work wear (arms, body and legs covered) and gloves TC: No TC available for this assessment	--	--
Critical Use: Ornamentals plants and trees			
Cutting, sorting, bundling, carrying Outdoor Work rate: 8 hours/day DT ₅₀ : 7 days DFR: 3µg/cm ² /kg a.s./ha Interval between applications: 7 days			
Application rate		3 x 1 kg.as/ha	
EFSA model 2014 (AOEM Excel calculator) Body weight: 60 kg	Potential TC: 14000 cm ² /person/h	0.8820	1102.5
	Work wear (arms, body and legs covered) TC: 5000 cm ² /person/h	0.3150	394
	Work wear (arms, body and legs covered) and gloves TC: 1400 cm ² /person/h	0.0882	110

Conclusion for outdoor uses:

There is no unacceptable risk anticipated for worker re-entering in treated crop **except for ornamentals, grapes, orchards.**

Indoor:

		Copper under the form of Bordeaux mixture	
Model data	Level of PPE	Total absorbed dose (mg/kg bw/day)	% of systemic AOEL
Critical use: fruiting vegetables Covering: tomato, eggplant, pepper, zucchini, gherkin, melon, watermelon, pumpkin, cucumber, lettuce, spinach			
Reaching, picking Indoor Work rate: 8 hours/day, DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days			
Number of applications and application rate		4 x 1 kg.as/ha (as worst case)	
EFSA model 2014 (AOEM Excel calculator) Body weight: 60 kg	Potential TC: 5800 cm ² /person/h	0,3915	489
	Work wear (arms, body and legs covered) TC: 2500 cm ² /person/h	0,1688	211
	Work wear (arms, body and legs covered) and gloves TC: 580 cm ² /person/h	0,0392	49
Critical Use: Strawberry			
Reaching, picking Indoor Work rate: 8 hours/day, DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days			
Number of applications and application rate		4 x 1 kg. as/ha	
EFSA model 2014 (AOEM Excel calculator) Body weight: 60 kg	Potential TC: 5800 cm ² /person/h	0,3915	489
	Work wear (arms, body and legs covered) TC: 3000 cm ² /person/h	0,2025	253
	Work wear (arms, body and legs covered) and gloves TC: 750 cm ² /person/h	0,0506	63
Critical Use: Ornamentals			
Cutting, sorting, bundling, carrying Indoor			

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Work rate: 8 hours/day DT ₅₀ : 7 days DFR: 3µg/cm ² /kg a.s./ha Interval between applications: 7 days			
Application rate		3 x 1 kg.as/ha	
EFSA model 2014 (AOEM Excel calculator) Body weight: 60 kg	Potential TC: 14000 cm ² /person/h	0.08953	1119
	Work wear (arms, body and legs covered) TC: 5000 cm ² /person/h	0.3283	410
	Work wear (arms, body and legs covered) and gloves TC: 1400 cm ² /person/h	0.1015	127

Conclusion for indoor uses:

There is no unacceptable risk anticipated for worker re-entering in treated crop **except for ornamentals**.

For details of personal protective equipment for workers, refer to the Decision in Appendix 1.

3.4.4 Bystander exposure

Consideration of acute exposure should only be made where an AAOEL has been established during an approval, review or renewal evaluation of an active substance, i.e. no acute operator or bystander exposure assessments can be performed with the AOEM model where no AAOEL has been set¹³.

According to EFSA Guidance on the assessment of exposure of operators, workers, residents and bystanders in risk assessment for plant protection products (EFSA Journal 2014;12(10):3874): “No bystander risk assessment is required for PPPs that do not have significant acute toxicity or the potential to exert toxic effects after a single exposure. Exposure in this case will be determined by average exposure over a longer duration, and higher exposures on one day will tend to be offset by lower exposures on other days. Therefore, exposure assessment for residents also covers bystander exposure.”

No AAOEL has been set for copper. Thus, for this active substance, residents exposure assessment covers bystanders exposure.

3.4.5 Resident exposure

Outdoor: Residential exposure was assessed according to EFSA model incorporating a distance of 10 meters for high crops and 2-3 meters for low crops. An acceptable risk was determined for residents (adult and child) without drift reduction technology.

		Copper under the form of Bordeaux mixture	
Model data		Total absorbed dose (mg/kg bw/day)	% of systemic AOEL
Apricot (covers other orchards)			
AOEM calculator (EFSA Model) Tractor mounted, upward application** Buffer zone: 10 m			

¹³ Guidance on the assessment of exposure of operators, workers, residents and bystanders in risk assessment for plant protection products (SANTE-10832-2015 rev. 1.7, 2017)

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Drift reduction technology: no DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days Volume min: 400 L/ha			
Number of applications and application rate		4 x 1 kg as./ha (5 x 0.8 kg as./ha*)	
Resident child Body weight: 10 kg	Sum (mean)	0,0481 (0,0394)	60 (49)
Resident adult Body weight: 60 kg	Sum (mean)	0,0255 (0,0208)	32 (26)
Grapes			
AOEM calculator (EFSA Model) Tractor mounted, upward application** Buffer zone: 10 m Drift reduction technology: no DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days Volume min: 400 L/ha			
Number of applications and application rate		4 x 1 kg a.s./ha (covers 5 x 0.8 kg.a.s./ha)	
Resident child Body weight: 10 kg	Sum (mean)	0.0450	56
Resident adult Body weight: 60 kg	Sum (mean)	0.0244	30.5
Ornamental plants and trees (trees as worst case)			
AOEM calculator (EFSA Model) Tractor mounted, upward application** Buffer zone: 10 m Drift reduction technology: no DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days Volume min: 400 L/ha			
Number of applications and application rate		3 x 1 kg a.s./ha	
Resident child Body weight: 10 kg	Sum (mean)	0.0436	54.5
Resident adult Body weight: 60 kg	Sum (mean)	0.0236	29.5
Cucumber Covering: tomato, eggplant, pepper, artichoke, zucchini, gherkin, melon, watermelon, pumpkin, leek, cardoon, lettuce, spinach, broccoli, cauliflower, onion, garlic, shallot, bean, pea, lentils, strawberry.			
AOEM calculator (EFSA Model) Tractor mounted, downward application Buffer zone: 2-3 m Drift reduction technology: no DT ₅₀ : 7 days DFR: 3 µg/cm ² /kg a.s./ha Interval between treatments: 7 days Volume min: 400 L/ha			

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Number of applications and application rate		4 x 1 kg a.s./ha (5 x 0.8 kg.sa/ha)	
Resident child Body weight: 10 kg	Sum (mean)	0.0295 (0.0245)	37 (31)
Resident adult Body weight: 60 kg	Sum (mean)	0.0144 (0.0119)	18 (15)

* covers the orchards for which a higher volume of water is claimed in GAP table by the applicant

** covers manual application for grape, olive or ornamental plants and trees

Indoor: In the context of indoor uses, resident exposure is not relevant.

Conclusion:

In conclusion, according to the EFSA model, there is **no unacceptable** risk anticipated for resident (adult and child) after incidental exposure to copper.

3.4.6 Combined exposure

Not relevant. The product contains only one active substance.

3.5 Residues and consumer exposure (Part B, Section 7)

As ornamental plants and trees are non edible commodities, the respective intended uses were not assessed.

For France, an exceedance of the current MRL for copper as laid down in Reg. (EC) 396/2005 of 50 mg/kg in grapes, 5 mg/kg in pome fruits, 5 mg/kg in stone fruits, 30 mg/kg in tree nuts and pistachio, 30 mg/kg in olive, 20 mg/kg in citrus trees, 5.0 mg/kg in outdoor and indoor tomato and aubergine, 5.0 mg/kg in outdoor and indoor edible peel cucurbits, 5.0 mg/kg in onion, garlic and shallot, 5.0 mg/kg in other roots and tuber vegetables except sugarbeet (ie.: parsnip, Jerusalem artichoke, parsley, celeriac, horseradish and scorzonera), 100 mg/kg in indoor lettuce and other salads, 20 mg/kg in fresh beans and peas with pods and fresh beans without pods is not expected.

Due to MRL exceedance, the uses on strawberries (outdoor and indoor), inedible peel cucurbits (outdoor and indoor), spring onion, spinaches and similar leaves (outdoor and indoor), pepper (outdoor and indoor) and potatoes cannot be recommended.

Due to insufficient residue trials, the uses on flowering brassica, lettuce (outdoor), fresh peas without pods, fresh lentils, carrots, beetroot, leeks, celery, cardoon, and artichokes cannot be recommended.

The acute exposure calculations were not carried out because an acute reference dose (ARfD) was not deemed necessary for copper.

For chronic intake of copper residues, the calculation includes uncertainties linked to the methodology. Therefore, zRMS considers that the risk assessment for consumers cannot be finalised

zRMS considers no firm conclusion can be reached for any of the intended uses of the product SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80).

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Summary for SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80)

Crop	PHI for SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) requested by applicant	PHI/ Withholding period* sufficiently supported for Copper	PHI for SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) proposed by zRMS	zRMS Comments (if different PHI proposed)
<i>Outdoor uses</i>				
Citrus trees	14 days	Yes	14 days	
Stone fruits	F**	Yes	F	
Pome fruits	F	Yes	F	
Tree nuts, pistachio	F	Yes	F	
Olive	14 days	Yes	14 days	
Grape	21 days	Yes	21 days	
Tomato, aubergine	10 days	Yes	10 days	
Pepper	14 days	n.a. (MRL exceedance)	-	Not recommended use
Leek, artichoke	7 days	No	-	Not recommended use
Spring onion	7 days	n.a. (MRL exceedance)	-	Not recommended use
Celery, cardoon	7 days	No	-	Not recommended use
Flowering brassica	14 days	No	-	Not recommended use
Cucurbits edible peel	3 days	Yes	3 days	
Cucurbits inedible peel	7 days	n.a. (MRL exceedance)	-	Not recommended use
Onion	3 days	Yes	3 days	
Garlic, shallot	3 days	Yes	3 days	
Lettuce	7 days	No	-	Not recommended use
Other salad plants including leaves and sprouts of brassicacea	7 days	No	-	Not recommended use
Spinach and similar (leaves)	7 days	n.a. (MRL exceedance)	-	Not recommended use
Fresh beans/peas with pods	3 days	Yes	3 days	
Fresh beans without pod	3 days	Yes	3 days	
Fresh peas without pods, fresh lentils	3 days	No	-	Not recommended use
Potato	14 days	n.a. (MRL exceedance)	-	Not recommended use

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Crop	PHI for SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) requested by applicant	PHI/ Withholding period* sufficiently supported for Copper	PHI for SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) proposed by zRMS	zRMS Comments (if different PHI proposed)
Carrot	14 days	No	-	Not recommended use
Other roots and tuber vegetables except sugarbeet (parsnip, Jerusalem artichoke, parsley, celeriac, horseradish and scorzonera)	14 days	Yes	14 days	
Beetroot	14 days	No	-	Not recommended use
Strawberry	7 days	n.a. (MRL exceedance)	-	Not recommended use
Ornamental plant and trees	NR		NR	Not assessed (non edible commodity)
Indoor uses				
Tomato, aubergine	3 days	Yes	3 days	
Pepper	14 days	n.a. (MRL exceedance)	-	Not recommended use
Cucurbits edible peel	3 days	Yes	3 days	
Cucurbits inedible peel	7 days	n.a. (MRL exceedance)	-	Not recommended use
Lettuce and other salad plants including leaves and sprouts of brassicacea	7 days	Yes	7 days	
Spinach and similar (leaves)	7 days	n.a. (MRL exceedance)	-	Not recommended use
Strawberry	7 days	n.a. (MRL exceedance)	-	Not recommended use
Ornamental plant and trees	NR		NR	Not assessed (non edible commodity)

NR: not relevant

* Purpose of withholding period to be specified

** F: PHI is defined by the application stage at last treatment (time elapsing between last treatment and harvest of the crop).

3.6 Environmental fate and behaviour (Part B, Section 8)

The fate and behaviour in the environment have been evaluated according to the requirements of Regulation (EC) No 1107/2009. Appropriate endpoints from the EU conclusions were used to calculate predicted environmental concentration (PEC) values for the active substance for the intended use patterns. In cases where deviations from the EU agreed end-points were considered appropriate (for example when additional studies are provided), such deviations were highlighted and justified accordingly.

The PEC of copper in soil, surface water and groundwater have been assessed according to FOCUS guidance documents, with standard FOCUS scenarios to obtain outputs from the FOCUS models, and the endpoints established in the EU conclusions or agreed in the assessment based on new data provided.

PECsoil and PECsw, accumulation (derived for 10 years) can be used for the risk assessment for the non-target terrestrial organisms for all intended uses.

Given the uncertainties identified by zRMS in the notifier's exposure calculation (FOCUS STEP 1-2 for all entries to water bodies and FOCUS STEP 1-2 PECsw including mitigation measures) and the absence of results for all FOCUS scenarios, PECsw derived for the active substance cannot be used for the ecotoxicological risk assessment. As a consequence, the risk assessment cannot be finalised for the non-target aquatic organisms for all intended uses.

PECgw for copper do not occur at levels exceeding those mentioned in Directive 98/83/CE¹⁴. Therefore, no unacceptable risk of groundwater contamination is expected for all intended uses.

Based on vapour pressure, no significant contamination of the air compartment is expected for the intended uses.

3.7 Ecotoxicology (Part B, Section 9)

The ecotoxicological risk assessment of the formulation was performed according to the requirements of Regulation (EC) No 1107/2009. Appropriate endpoints from the EU conclusions for the active substance were used for the intended use patterns. In cases where deviations from the EU agreed endpoints were considered appropriate (for example when additional studies are provided), such deviations were highlighted and justified accordingly.

An EFSA' Statement of the PPR panel on a framework for conducting the environmental exposure and risk assessment for transition metals when used as active substances in plant protection products was recently published (2021). This document provides useful recommendations upon applicability of new methodologies in the context of transition metals and possible areas of development for assessing the risk from transition metals used in PPPs. However, it does not provide valid tools for exposure assessment in the environment and toxicity estimation upon non-target organisms. Furthermore, no clear specific risk assessment schemes for transition metals used as active substances in PPPs is provided. Therefore, the risk assessment and conclusion are based on the methodology agreed by the experts during the renewal approval of the active substance. The EU-agreed endpoints recommended in the EFSA journal (EFSA Journal 2018;16(1):5152) were considered for the Art. 43 dossiers for copper compounds.

Based on the guidance documents, the risks for **non-target arthropods other than bee** and non-target **terrestrial plants** are acceptable for the intended uses.

For birds and mammals, the risk is not acceptable at Tier 1 for all intended uses. The arguments provided by the applicant to refine the risk assessment are identical to those that were considered insufficient at the European level. Therefore, without further data, the risk assessment for birds and mammals cannot be finalised except for applications under permanent greenhouse.

For aquatic organisms, as the toxicity reference value for copper proposed by the applicant was based on an approach rejected at European level, it could not be used. In addition, no reliable PECsed were provided by the applicant for all uses. Therefore, the risk assessment for non-target aquatic species could not be finalised for uses in open field, in tunnels or in permanent greenhouses with soil-bound cultivation. For

¹⁴ Council Directive 98/83/EC of 3 November 1998 on the quality of water intended for human consumption

uses in permanent greenhouses with soil-less cultivation, the exposure of aquatic organisms to the active substance from the use of the product SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) is considered negligible.

For bees, the risk assessment provided by the applicant is based on the EFSA Guidance Document¹⁵.

For adult honey bees, the risk is not acceptable at Tier 1 for all intended uses. Higher-tier studies (cage and tunnel tests) are available and demonstrate that no adverse effects on adult honey bees are expected for all intended uses.

For honey bee larvae, the risks are not acceptable at Tier 1 for all intended uses except for olives, hazelnuts and pistachios, tomatoes, lettuce and potatoes, in open-field or in walk-in tunnels. For all other uses in open-field or in walk-in tunnels, the higher-tier studies are not sufficient to demonstrate the absence of adverse effects of the product SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) on honey bee larvae. Therefore, the risk assessment for honey bee larvae cannot be finalised for applications in open-field or in walk-in tunnels, for all requested uses except for olives, hazelnuts and pistachios, tomatoes, lettuce and potatoes.

For bumble bees, no acute risk assessment was provided by the applicant, although standard study protocols are available. Therefore, the risk assessment for bumble bees cannot be finalised for applications in open-field or in walk-in tunnels.

Overall, the risk for bees cannot be finalised for all requested uses except for applications under permanent greenhouse. For these structures, the following precautionary statement should be applied: "May affect pollinators. Avoid unnecessary exposure".

For soil organisms, since no reliable PEC soil are available, a Tier 1 risk assessment cannot be conducted. For earthworms, the higher tier earthworm field trial data from a study conducted over 10 years with copper application every year demonstrates that there is an acceptable risk to earthworms for applications up to 4 kg Cu/ha/year. Therefore, an acceptable risk for earthworms is demonstrated for all intended uses of SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80).

For other soil meso- and macro-organisms, no higher-tier studies are available and extrapolating the results of the multiyear field study with earthworms to other soil meso- and macro-organisms was not supported by the experts at the Peer Review experts' meeting 169. Therefore, the risk for soil macro-organisms other than earthworms could not be finalised for all intended uses except for uses under permanent greenhouse with soil-less cultivation.

For soil micro-organisms, based on a lack of effect at field level, the risks to soil micro-organisms are acceptable for the intended uses.

3.8 Relevance of metabolites (Part B, Section 10)

An assessment was conducted according to the SANCO/221/2000 guidance document. Please refer to environmental fate and behaviour above for conclusion on the risk of groundwater contamination.

4 Conclusion of the national comparative assessment (Art. 50 of Regulation (EC) No 1107/2009)

SUPER BOUILLIE MACCLESFIELD 80 (Super Bouillie Macclesfield 80) contains copper compounds, which is approved as a candidate for substitution because it fulfills PBT criteria (Persistent and Toxic);

Steps 1 and 2 (French guidance document 27 July 2015):

- **Taking into account the agronomic interest, especially in the context of organic farming**

¹⁵ EFSA Guidance Document on the risk assessment of plant protection products on bees (*Apis mellifera*, *Bombus* spp. and solitary bees) EFSA Journal 2013;11(7):3295

In accordance with Article 50, paragraphs 1.b) 1.c) and 1.d) of Regulation (EC) N°1107/2009,

- considering the absence of plant protection products or non-chemical methods of prevention or control allowing to consider a substitution of the product without major practical or economic disadvantage, and especially in the frame of organic farming,
- considering also the need to guarantee a diversity of the modes of action to reduce the emergence of resistance in target microorganisms,
- considering the need to take into account the minor uses of the product,

the substitution of the product will not be considered for all intended uses.

5 Further information to permit a decision to be made or to support a review of the conditions and restrictions associated with the authorisation

When the conclusions of the assessment is “Not acceptable”, please refer to relevant summary under point 3, “Background of authorisation decision and risk management”.

5.1.1 Post-authorisation monitoring

N.A

5.1.2 Post-authorisation data requirements

None.

Appendix 1 Copy of the product authorisation DAMM

Docusign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Décision relative à une demande de renouvellement de l'autorisation de mise sur le marché d'un produit phytopharmaceutique

Vu les dispositions du règlement (CE) n° 1107/2009 du 21 octobre 2009 et de ses textes d'application,

Vu le code rural et de la pêche maritime, notamment le chapitre III du titre V du livre II des parties législative et réglementaire,

*Vu la demande de renouvellement de l'autorisation de mise sur le marché, suite au renouvellement de l'approbation de la substance active composés du cuivre, du produit phytopharmaceutique **SUPER BOUILLIE MACCLESFIELD 80***

de la société **INDUSTRIAS QUIMICAS DEL VALLES**

enregistrée sous le **n° 2019-3931**

Vu les conclusions de l'évaluation de l'Anses du 24 juin 2022,

Considérant que les informations disponibles ne permettent pas de s'assurer de la performance des équipements de protection individuelle pour cette formulation de type poudre mouillable,

Considérant en conséquence qu'un risque d'effet nocif pour les opérateurs et les travailleurs lié à l'utilisation du produit ne peut être exclu,

Considérant qu'en conséquence, les exigences mentionnées à l'article 29 du règlement (CE) n° 1107/2009 ne sont plus remplies,

L'autorisation de mise sur le marché du produit phytopharmaceutique désigné ci-après **n'est pas renouvelée** en France.

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Informations générales sur le produit	
Nom du produit	SUPER BOUILLIE MACCLESFIELD 80
Type de produit	Produit de référence
Titulaire	INDUSTRIAS QUIMICAS DEL VALLES Av. Rafael Casanova 81 08100 MOLLET DEL VALLES Espagne
Formulation	Poudre mouillable (WP)
Contenant	200 g/kg – cuivre (sous forme de 733,4 g/kg de bouillie bordelaise)
Numéro d'intrant	7500716
Numéro d'AMM	7500716
Fonction	Fongicide et bactéricide
Gamme d'usage	Professionnel

A Maisons-Alfort, le 15/07/2025

DocuSigned by:
Charlotte Grastilleur

AE281A955A42454...

Directrice générale déléguée
en charge du pôle produits réglementés
Agence nationale de sécurité sanitaire de
l'alimentation, de l'environnement et du travail (ANSES)

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



ANNEXE : Conditions de mise sur le marché demandées

Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
12053301 Agrumes*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12053204 Agrumes * Trt Part.Aer * Chancre du collet	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12053200 Agrumes*Trt Part.Aer.*Maladies des feuilles et fruits	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12103201 Amandier*Trt Part.Aer.*Cloque(s)	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12103202 Amandier*Trt Part.Aer.*Coryneum	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12103203 Amandier*Trt Part.Aer.*Moniliose(s)	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

Docusign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
14153204 Arbres et arbustes*Trt Part.Aer.*Anthracnose(s)	5 kg/ha	3/an	Non applicable
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
00002019 Arbres et arbustes*Trt Part.Aer.*Maladies des taches foliaires	5 kg/ha	3/an	Non applicable
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
00002023 Arbres et arbustes*Trt Part.Aer.*Moniliose(s)	5 kg/ha	3/an	Non applicable
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
16103301 Artichaut*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
16103204 Artichaut*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
16103202 Artichaut*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
16173203 Betterave potagère*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs, ni de déterminer l'efficacité du produit.		

SUPER BOUILLIE MACCLESFIELD 80

AMM n° 7500716

Page 4 sur 17

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
01108014 Carotte*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque de dépassement des limites maximales de résidus sur carotte ni d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
16203204 Carotte*Trt Part.Aer.*Champignons (pythiacées)	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque de dépassement des limites maximales de résidus sur carotte ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
16203203 Carotte*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque de dépassement des limites maximales de résidus sur carotte ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
16253301 Céleris*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car transitoire et transformé en "01109011 Céleri-branche*Trt.Part.Aer.*Bactériose(s)" et "01109011 Carotte*Trt.Part.Aer.*Bactériose(s)".		
16253201 Céleris*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car transitoire et transformé en "19273201 Céleri-branche*Trt.Part.Aer.*Maladies des taches brunes" et en "16203203 Carotte*Trt.Part.Aer.*Maladies des taches brunes".		
16253204 Céleris * Trt Part.Aer.* Mildiou(s)	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car transitoire et transformé en "01109010 Céleri-branche*Trt.Part.Aer.*Mildiou(s)" et "16203204 Carotte*Trt.Part.Aer.*Champignons (pythiacées)".		
01109011 Céleri-branche*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
19273201 Céleri-branche*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
01109010 Céleri-branche* Trt. Part. Aer. * Miliou(s)".	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
12203201 Cerisier*Trt Part.Aer.*Anthracnose(s)	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12203208 Cerisier*Trt Part.Aer.*Moniliose(s) et pourriture grise	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12253203 Chataignier*Trt Part.Aer.*Septoriose(s)	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
00516023 Choux à inflorescence*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
00516026 Choux à inflorescence*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
00516027 Choux à inflorescence*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
01116017 Cucurbitacées à peau comestible*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
16323205 Cucurbitacées à peau comestible*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
16323204 Cucurbitacées à peau comestible*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
16753201 Cucurbitacées à peau non comestible*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
16753208 Cucurbitacées à peau non comestible*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
17053201 Cultures florales et plantes vertes*Trt Part.Aer.*Maladies des taches foliaires	5 kg/ha	3/an	Non applicable
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
17403200 Cultures florales et plantes vertes*Trt Part.Aer.*Maladies diverses	5 kg/ha	3/an	Non applicable
Motivation du refus : L'usage est refusé car transitoire et transformé en " 17053201 - Cultures florales et plantes vertes*Trt Part. Aer.* Maladies des taches foliaires" et en " 17403204 - Cultures florales et plantes vertes* Trt Part. Aer.* Mildiou(s)".			
17403204 Cultures florales et plantes vertes*Trt Part.Aer.*Mildiou(s)	5 kg/ha	3/an	Non applicable
Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.			
17403203 Cultures florales et plantes vertes*Trt Part.Aer.*Rouille(s)	5 kg/ha	3/an	Non applicable
Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs ni de déterminer l'efficacité du produit.			
16503201 Epinard*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	7
Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.			
16503204 Epinard*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	7
Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.			
16553207 Fraisier*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	7
Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.			
16553203 Fraisier*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	7
Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.			

SUPER BOUILLIE MACCLESFIELD 80

AMM n° 7500716

Page 8 sur 17

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
Fruits à coque*Trt Part.Aer.*Alternariose(s)	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12603303 Fruits à pépins*Trt Part.Aer.*Feu bactérien	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12603205 Fruits à pépins*Trt Part.Aer.*Moniliose(s)	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
16573301 Haricots et Pois écosés frais*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
16573202 Haricots et Pois écosés frais*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs. L'usage sur pois écosé frais est refusé car les données disponibles ne permettent pas de vérifier le respect des limites maximales de résidus de la substance active.		
16573206 Haricots et Pois écosés frais*Trt Part.Aer.*Rouille(s)	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs ni de déterminer l'efficacité du produit. L'usage sur pois écosé frais est refusé car les données disponibles ne permettent pas de vérifier le respect des limites maximales de résidus de la substance active.		

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
00516012 Haricots et pois non écosés frais*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
00516013 Haricots et pois non écosés frais*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
00516020 Haricots et pois non écosés frais*Trt Part.Aer.*Rouille(s)	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs ni de déterminer l'efficacité du produit.		
16703208 Laitue*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque de dépassement des limites maximales de résidus en plein champ ni d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
16603207 Laitue*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque de dépassement des limites maximales de résidus en plein champ ni d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
16603201 Laitue*Trt Part.Aer.*Pourriture grise et sclérotinioses	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque de dépassement des limites maximales de résidus en plein champ ni d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, ni de déterminer l'efficacité du produit.		
00211018 Noisetier*Trt Part.Aer.*Dépérissement cryptogamique	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		

SUPER BOUILLIE MACCLESFIELD 80

AMM n° 7500716

Page 10 sur 17

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
16423301 Oignon*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque d'effet nocif pour les opérateurs et les travailleurs.		
16053205 Oignon*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque d'effet nocif pour les opérateurs et les travailleurs.		
16803201 Oignon*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	3
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque d'effet nocif pour les opérateurs et les travailleurs.		
12503201 Olivier*Trt Part.Aer.*Maladies des fruits	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs. Cet usage intègre l'usage revendiqué "Olivier*Trt Part.Aer.*Colletotrichum gloeosporioides".		
12553230 Pêcher - Abricotier *Trt Part.Aer.*Chancres à champignons	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12553233 Pêcher - Abricotier*Trt Part.Aer.*Moniliose(s)	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12853201 Pistachier*Trt Part.Aer.*Maladies du feuillage	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs. Cet usage intègre l'usage revendiqué "Pistache*Trt.Part.Aer.*Alternaria spp".		

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés			
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
16843201 Poireau*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	7
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
01140015 Poivron*Trt Part.Aer.*Bactériose(s)	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
16863205 Poivron*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
16863204 Poivron*Trt Part.Aer.*Mildiou(s)	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
15653202 Pomme de terre*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	14
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs.		
12653203 Prunier*Trt Part.Aer.*Cloque(s)	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
12653206 Prunier*Trt Part.Aer.*Coryneum et polystigma	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages refusés

Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)
12653204 Prunier*Trt Part.Aer.*Moniliose(s)	5 kg/ha	4/an	F
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
17303205 Rosier*Trt Part.Aer.*Mildiou(s)	5 kg/ha	3/an	Non applicable
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		
Rosier*Trt Part Aer.*Monilia spp.	5 kg/ha	3/an	Non applicable
	Motivation du refus : L'usage est refusé car agronomiquement non pertinent.		
16953207 Tomate - Aubergine*Trt Part.Aer.*Maladies des taches brunes	5 kg/ha	4/an	10
	Motivation du refus : L'usage est refusé car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs.		

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages retirés					
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)	Délai accordé pour la vente et la distribution	Délai accordé pour le stockage et l'utilisation des stocks
12203301 Cerisier*Trt Part.Aer.*Bactériose(s)	1,25 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12203204 Cerisier*Trt Part.Aer.*Coryneum et polystigma	1,25 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
16753301 Cucurbitacées à peau non comestible*Trt Part.Aer.*Bactériose(s)	-	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus et ni un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
16553301 Fraisier*Trt Part.Aer.*Bactériose(s)	12,5 kg/ha	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus ni un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12453301 Fruits à coque*Trt Part.Aer.*Bactériose(s)	1,25 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées. Cet usage intègre les usages revendiqués : "Pistache*Trt.Part.Aer.*Xanthomonas spp." et "Pistache*Trt.Part.Aer.*Pseudomonas spp.".					

SUPER BOUILLIE MACCLESFIELD 80

AMM n° 7500716

Page 14 sur 17

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages retirés					
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)	Délai accordé pour la vente et la distribution	Délai accordé pour le stockage et l'utilisation des stocks
12603301 Fruits à pépins*Trt Part.Aer.*Bactériose(s)	1,25 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12603201 Fruits à pépins*Trt Part.Aer.*Chancre européen	2,5 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12603203 Fruits à pépins*Trt Part.Aer.*Tavelure(s)	1,25 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
16613301 Laitue*Trt Part.Aer.*Bactériose(s)	12,5 kg/ha	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque de dépassement des limites maximales de résidus en plein champ ni d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12503301 Olivier*Trt Part.Aer.*Bactériose(s)	1,25 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12503203 Olivier*Trt Part.Aer.*Maladies du feuillage	1,25 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages retirés					
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)	Délai accordé pour la vente et la distribution	Délai accordé pour le stockage et l'utilisation des stocks
12553303 Pêcher - Abricotier*Trt Part.Aer.*Bactériose(s)	0,625 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12553203 Pêcher - Abricotier*Trt Part.Aer.*Cloque(s)	2,5 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12553232 Pêcher - Abricotier*Trt Part.Aer.*Coryneum et polystigma	1,25 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
15653201 Pomme de terre*Trt Part.Aer.*Mildiou(s)	25 kg/ha	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure d'un risque de dépassement des limites maximales de résidus, aux conditions d'emploi revendiquées.					
01801019 PPAMC*Trt Part.Aer.*Bactériose(s)	12,5 kg/ha	3/an	21	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car il n'a pas été soutenu lors de la demande de renouvellement de l'autorisation du produit phytopharmaceutique.					
19993200 PPAMC*Trt Part.Aer.*Maladies fongiques	6,25 kg/ha	3/an	21	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car il n'a pas été soutenu lors de la demande de renouvellement de l'autorisation du produit phytopharmaceutique.					

SUPER BOUILLIE MACCLESFIELD 80

AMM n° 7500716

Page 16 sur 17

Super Bouillie Macclesfield 80
Part A - National Assessment
FRANCE DEPR version

DocuSign Envelope ID: D9B29F25-0FEA-460B-9566-2A35A0C47B19



Liste des usages retirés					
Usages	Dose d'emploi	Nombre maximum d'applications	Délai avant récolte (jours)	Délai accordé pour la vente et la distribution	Délai accordé pour le stockage et l'utilisation des stocks
12653301 Prunier*Trt Part.Aer.*Bactériose(s)	1,25 kg/hL	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
16953301 Tomate - Aubergine*Trt Part.Aer.*Bactériose(s)	20 kg/ha	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
16953201 Tomate - Aubergine*Trt Part.Aer.*Mildiou(s)	6,25 kg/ha	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12703301 Vigne*Trt Part.Aer.*Bactériose(s)	20 kg/ha	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					
12703203 Vigne*Trt Part.Aer.*Mildiou(s)	15 kg/ha	-	-	6 mois à compter de la présente décision	18 mois à compter de la présente décision
Motivation du retrait : L'usage est retiré car les données disponibles ne permettent pas d'exclure un risque d'effet nocif pour les opérateurs et les travailleurs, aux conditions d'emploi revendiquées.					

Appendix 2 Copy of the product label

The draft product label as proposed by the applicant is reported below. The draft label may be corrected with consideration of any new element. The label shall reflect the detailed conditions stipulated in the Decision.

Mars 2019
Page 1 / 10

SUPER BOUILLIE MACCLESFIELD 80

FONGICIDE ET BACTÉRICIDE

Fongicide et bactéricide préventif à large spectre pour les vergers, la vigne, les cultures légumières, la pomme de terre, la betterave sucrière, les fraisiers et les cultures ornementales

Autorisation de Mise sur le Marché (A.M.M.) N° 7500716

Détenteur de l'A.M.M. : Industrias Químicas del Vallés, S.A. (Av. Rafael Casanova 81 - 08100 Mollet del Vallés - Barcelona, Espagne)

Contient 200 g/kg (20.0% p/p) de cuivre (sous forme de bouillie bordelaise)
Poudre mouillable (WP)

RESERVÉ À UN USAGE EXCLUSIVEMENT PROFESSIONNEL.

SUPER BOUILLIE MACCLESFIELD 80 (A.M.M. N° 7500716)		
Attention 	H410	Très toxique pour les organismes aquatiques, entraîne des effets néfastes à long terme.
	P391	Recueillir le produit répandu.
	P501	Éliminer le contenu/récipient conformément à la réglementation locale.
	Délai de rentrée sur la parcelle : 6 heures et, en cas d'application en milieu fermé, de 8 heures.	
	SPe3	Pour protéger les organismes aquatiques, respecter une zone non traitée par rapport aux points d'eau de : - 5 mètres avec dispositif végétalisé permanent de 5 mètres pour les usages sur agrumes, olive, vigne, tomate, aubergine, poivron, piment, poireau, oignon de printemps, ciboule, artichaut, céleri, cardon, brocoli, chou-fleur, concombre, courgette, cornichon, melon, pastèque, potiron, oignon, ail, échalote, laitue et autres salades, épinard, pourpier, feuilles de bettes, haricots non écosés et écosés frais, pois non écosés et écosés frais, lentilles fraîches, pomme de terre, betterave potagère, carotte et autres légumes racines, betterave sucrière et fraisier. - 20 mètres avec dispositif végétalisé permanent de 5 mètres pour les usages sur fruits à noyau, fruits à pépins et fruits à coque. - 20 mètres en plein champ pour les usages sur cultures ornementales.
EUH401	Respectez les instructions d'utilisation afin d'éviter les risques pour la santé humaine et l'environnement.	
SP1	Ne pas polluer l'eau avec le produit ou son emballage. Ne pas nettoyer le matériel d'application près des eaux de surface. Éviter la contamination via les systèmes d'évacuation des eaux à partir des cours de ferme ou des routes.	
FABRICANT :		
Adresse Tél. : xxx.xxx.xxx.xxx		

EN CAS D'URGENCE

Composer le 15, le 112 ou contacter le centre anti-poison le plus proche

Puis signaler vos symptômes au réseau PhytAttitude, N° Vert : 0 800 887 887 (appel gratuit depuis un poste fixe).



Fiche de Données de Sécurité disponible sur demande pour les professionnels (ou sur www.quickfds.com).

Lire les instructions ci-jointes avant emploi.

EMB :

Mars 2019
Page 2 / 10

N° de lot et date de fabrication : voir emballage
Version 1.0
XX kg e

Mars 2019
Page 3 / 10

PREMIERS SOINS

En cas d'inhalation :

Transporter la victime à l'air libre et la garder au chaud et au repos.

En cas de contact avec les yeux :

Laver abondamment avec de l'eau douce et propre durant au minimum 15 minutes en maintenant les paupières écartées. Retirer les lentilles de contact si la victime en porte après 5 minutes de rinçage. Consulter un spécialiste.

En cas de contact avec la peau :

Enlever les vêtements imprégnés et laver soigneusement la peau avec de l'eau et du savon ou utiliser un nettoyant connu. Ne pas utiliser de solvants ou de diluants. En cas d'irritation, consulter un médecin.

En cas d'ingestion :

Rincer la bouche avec de l'eau et consulter un médecin ou un centre antipoison. Montrer l'étiquette. Garder au repos.

Ne pas faire vomir.

Antidote : EDTA, BAL ou pénicillamine. Consulter immédiatement un médecin et lui montrer l'étiquette.

Traitement symptomatique : un lavage gastrique avec une solution lacto-albumine peut être nécessaire.

RECOMMANDATIONS D'EMPLOI

IMPORTANT : Lire attentivement les instructions de cette section afin de garantir une utilisation sûre et efficace de ce produit.

MODE D'ACTION

Le cuivre appartient au groupe FRAC M01. C'est un fongicide et bactéricide multi-site inorganique.

TABLEAU DES USAGES

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Citron, Mandarine, Orange, Pamplemousse, Clémentine, Limette	Alternariose (<i>Alternaria alternata</i> f. sp. <i>Citri</i>) Bactérioses (<i>Pseudomonas syringae</i> pv. <i>syringae</i>) Chancre du collet (<i>Phytophthora</i> spp.)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (feuilles visibles mais n'ont pas encore atteint leur taille finale) à 89 (fruits mûrs)	1000-2000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Abricotier, Cerisier, Pêcher, Nectarinier, Prunier, Mirabellier, Jujubier	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>syringae</i>) Chancres à champignons (<i>Cytospora leucostoma</i>) Cloques (<i>Taphrina deformans</i>) Cronium (<i>Stigmella campophila</i>) Monilioses (<i>Monilia</i> spp.)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 95 (sénescence - 50% des feuilles sont décolorées ou tombées) à 53 (éclatement des bourgeons)	400-1000 L/ha	-	20 mètres avec DVP** de 5 mètres
Cerisier	Anthraxose (<i>Blumeriella jappa</i>)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 95 (50% des feuilles sont décolorées ou tombées) à 53 (éclatement des bourgeons)	400-1000 L/ha	-	20 mètres avec DVP** de 5 mètres
Pommier, Poirier, Cognassier, Néflier, Nashi, Pommette	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>populans</i>) Chancre européen (<i>Nectria galligena</i>) Feu bactérien (<i>Erwinia amylovora</i>) Monilioses (<i>Monilia</i> spp.) Tavelure (<i>Venturia</i> spp.)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 91 (début sénescence - fin de la croissance des rameaux, le bourgeon terminal est développé, les feuilles sont toujours vertes) à 53 (éclatement des bourgeons)	500-1000 L/ha	-	20 mètres avec DVP** de 5 mètres

Mars 2019
Page 4 / 10

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Amandier	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>syringae</i>) Cloques (<i>Taphrina deformans</i>) Coryneum (<i>Stigmella camponotila</i>) Monilioses (<i>Monilia</i> spp.)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 95 (50% des feuilles sont décolorées ou tombées) à 53 (éclatement des bourgeons)	500-1000 L/ha	-	20 mètres avec DVP** de 5 mètres
Noyer, Noisetier, Châtaignier, Pistachier	Alternariose (<i>Alternaria</i> spp.) Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Dépérissement cryptogamique (<i>Cytospora</i> spp.) Septoriose (<i>Mycosphaerella</i> spp.)	3.75 à 5 kg/ha	4 (intervalle de 7 jours entre appl.)	BBCH 95 (50% des feuilles sont décolorées ou tombées) à 53 (éclatement des bourgeons)	500-1000 L/ha	-	20 mètres avec DVP** de 5 mètres
Olivier	Anthraxose (<i>Colletotrichum gloeosporioides</i>) Bactérioses (<i>Pseudomonas savastanoi</i> pv. <i>savastanoi</i>) Maladie de l'œil de paon (<i>Sclerotinia oleaginea</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 10 (développement des feuilles) à 85 (début de la coloration du fruit)	600-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Vigne (raisin de cuve et raisin de table) (champ)	Anthraxose (<i>Elsinoe ampelina</i>) Bactérioses (<i>Xanthomonas ampelina</i>) Mildiou (<i>Plasmopara viticola</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (3 feuilles étalées) à 81 (début de la maturation) et 91 (après la vendange : acûtement du bois terminé)	400-1000 L/ha	21 jours	5 mètres avec DVP** de 5 mètres
Tomate, Aubergine (champ/ sous serre)	Bactérioses (<i>Pseudomonas syringae</i> et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria solani</i> , <i>Colletotrichum</i> spp.) Mildiou (<i>Phytophthora infestans</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (5 feuilles étalées sur la tige principale) à 89 (fruits mûrs)	400-1000 L/ha	10 jours (plein champ) / 3 jours (sous serre)	5 mètres avec DVP** de 5 mètres
Poivron, Piment (champ/ sous serre)	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>Syringae</i> et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria solani</i> , <i>Colletotrichum</i> spp.) Mildiou (<i>Phytophthora capsici</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (5 feuilles étalées sur la tige principale) à 89 (fruits mûrs)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Poireau, Oignon de printemps, Ciboule Artichaut, Céleris (branche et rave), Cardon (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria</i> spp., <i>Ascochyta</i> spp.) Mildiou (<i>Bremia lactucae</i> , <i>Phytophthora</i> spp., <i>Plasmopara crustosa</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 14 (4 feuilles visibles/ étalées) à 51 (le bulbe commence à s'allonger/ la pousse principale commence à sortir)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres

Mars 2019
Page 5 / 10

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Brocoli, Chou-fleur (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas campestris</i> pv. <i>campestris</i>) Maladies des taches brunes (<i>Alternaria brassicae</i> , <i>Colletotrichum</i> spp.) Mildiou (<i>Hyaloperonospora brassicae</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 14 (4 feuilles étalées) à 59 (premiers pétioles visibles, fleurs toujours fermées)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Concombre, Courgette, Cornichon (champ/ sous serre)	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>Lachrymans</i> et <i>Xanthomonas campestris</i> pv. <i>cucurbitae</i>) Maladies des taches brunes (<i>Alternaria cucumerina</i> , <i>Colletotrichum lagenarium</i>) Mildiou (<i>Pseudoperonospora cubensis</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 10 (cotylédons étalés) à 89 (maturation complète)	400-1000 L/ha	3 jours	5 mètres avec DVP** de 5 mètres
Melon, Pastèque, Potiron (champ/ sous serre)	Bactérioses (<i>Pseudomonas syringae</i> pv. <i>Lachrymans</i> et <i>Xanthomonas campestris</i> pv. <i>cucurbitae</i>) Maladies des taches brunes (<i>Alternaria cucumerina</i> , <i>Colletotrichum lagenarium</i>) Mildiou (<i>Pseudoperonospora cubensis</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 10 (cotylédons étalés) à 89 (maturation complète)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres
Oignon, Ail, Echalote (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria porri</i> , <i>Colletotrichum coccoodes</i> , <i>Fleospora alli</i>) Mildiou (<i>Peronospora destructor</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 14 (4 feuilles visibles) à 47 (feuilles fanées dans 10% des plantes)	400-1000 L/ha	3 jours	5 mètres avec DVP** de 5 mètres
Laitue et autres salades : Mâche Scarole (endive à larges feuilles) Cresson Cresson de terre Roquette, Rucola Moutarde brune Feuilles et pousses de Brassica spp. (champ/ sous serre)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas axonopodis</i> pv. <i>Vitians</i>) Mildiou (<i>Bremia lactucae</i>) Maladies des taches brunes (<i>Alternaria sonchii</i> , <i>Marssonina paratoniciana</i>) Sclerotiniose (<i>Sclerotinia minor</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 12 (2 feuilles étalées) à 49 (la taille typique de la masse foliaire est atteinte)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres
Epinaud, Pourpier, Feuilles de bettes (champ/ sous serre)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria</i> spp., <i>Colletotrichum spinaciae</i>) Mildiou (<i>Peronospora farinosa</i> f. sp. <i>Spinaciae</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 12 (2 feuilles étalées) à 49 (la taille typique de la masse foliaire est atteinte)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres

Mars 2019
Page 6 / 10

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Haricots non écosés frais (haricots verts, etc.) Haricots écosés frais (fèves, flageolets, etc.) Pois non écosés frais (pois mange-tout) Pois écosés frais (pois potagers, pois frais, pois chiches) Lentilles fraîches (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas axonopodis</i> pv. <i>phaseoli</i>) Maladies des taches brunes (<i>Colletotrichum lindemuthianum</i> , <i>Sclerotinia</i> spp.) Mildiou (<i>Peronospora viciae</i> , <i>Phytophthora phaseoli</i>) Rouilles (<i>Uromyces</i> spp.)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 14 (4 feuilles étalées ou 4 vrilles formées) à 89 (maturation complète)	400-1000 L/ha	3 jours	5 mètres avec DVP** de 5 mètres
Pomme de terre	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas</i> spp.) Maladies des taches brunes (<i>Alternaria solani</i> , <i>Colletotrichum coccodes</i>) Mildiou (<i>Phytophthora infestans</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (5*** feuille de la tige principale étalée) à 85 (baies de la première inflorescence ocre ou brunâtres)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Betterave potagère Carotte Raifort Topinambours Panais Persil à grosse racine Radis Salsifis Rutabagas Navets (champ)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas campestris</i> pv. <i>carotae</i>) Maladies des taches brunes (<i>Alternaria dauci</i> , <i>Cercospora carotae</i> , <i>Colletotrichum</i> spp.) Mildiou (<i>Plasmopara nivea</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 15 (5 feuilles étalées) à 47 (racines, tiges ou tubercules ont atteint 70% de leur taille finale)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Betterave sucrière	Maladies du feuillage (<i>Cercospora beticola</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 39 (fermeture des lignes: environ 90% des plantes des lignes adjacentes se touchent) à 49 (racine de la betterave à sa taille de récolte)	400-1000 L/ha	14 jours	5 mètres avec DVP** de 5 mètres
Fraisier (champ/ sous serre)	Bactérioses (<i>Pseudomonas</i> spp. et <i>Xanthomonas fragariae</i>) Maladies des taches brunes (<i>Colletotrichum fragariae</i> , <i>Mycosphaerella fragariae</i>) Mildiou (<i>Phytophthora fragariae</i>)	3.75 à 5 kg/ha	5 (intervalle de 7 jours entre appl.)	BBCH 13 (3 feuilles étalées) à 85 (les premières fraises ont atteint la couleur spécifique de la variété)	400-1000 L/ha	7 jours	5 mètres avec DVP** de 5 mètres

Culture	Cible	Dose homologuée	Nombre maximal de traitements par an*	Conditions d'emploi	Volume de bouillie conseillé L/ha min-max	Délai avant récolte (DAR)	Zone non traitée par rapport aux points d'eau
Arbres et Arbustes d'ornement Bulbes ornementaux Cultures florales et plantes vertes Rosier (champ/ sous serre)	Anthracnoses et maladies des taches foliaires (<i>Colletotrichum</i> spp., <i>Septoria</i> spp.) Mildiou (<i>Peronospora</i> spp.) Monilioses (<i>Monilia</i> spp.) Rouilles (<i>Puccinia</i> spp.)	3.75 à 5 kg/ha	3 (intervalle de 7 jours entre appl.)	Tous les stades	400-1000 L/ha	-	20 mètres en plein champ

* Seules les utilisations entraînant une application totale maximale de 28 kg de cuivre par hectare sur une période de 7 ans sont autorisées.

** DVP : Dispositif Végétalisé Permanent

LIMITES MAXIMALES DE RÉSIDUS (LMR)

Les LMR sont consultables à l'adresse suivante : <http://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/>

PRÉCONISATIONS D'EMPLOI

SUPER BOUILLIE MACCLESFIELD 80 est un fongicide et bactéricide préventif à large spectre. A base de cuivre, il est plus efficace contre les spores que contre les maladies déjà établies, il est donc recommandé de l'appliquer avant l'apparition de la maladie, ou dès que les premiers symptômes de la maladie sont visibles.

Dans des conditions très favorables au développement des maladies visées, il est recommandé d'alterner l'utilisation de SUPER BOUILLIE MACCLESFIELD 80 avec d'autres produits contenant des substances actives ayant un mode d'action différent.

PRÉPARATION DE LA BOUILLIE

Remplir à moitié la cuve avec de l'eau et mettre en marche l'agitation. Verser la quantité nécessaire de SUPER BOUILLIE MACCLESFIELD 80 dans la cuve du pulvérisateur. Remplir la cuve avec de l'eau au volume requis. Maintenir l'agitation durant toute la durée de l'application.

APPLICATION

Éviter tout risque de dérive de pulvérisation. Traiter par temps calme.

Ne pas laisser la bouillie dans la cuve du pulvérisateur pendant de longues périodes par exemple le temps des repas.

Appliquer en pulvérisation foliaire. Pour assurer une efficacité optimale, le matériel de pulvérisation doit être calibré afin de bien appliquer le produit sur l'ensemble des parties de la culture à protéger et assurer une distribution uniforme du produit sur la culture. Ajuster le volume de bouillie en fonction du matériel utilisé et du volume de végétation afin d'assurer une couverture complète de la végétation à protéger.

Appliquer préventivement avec un intervalle recommandé de 7 jours entre les applications. Cependant, en règle générale, les applications doivent toujours être adaptées à l'évolution de la maladie, de la culture traitée et aux conditions climatiques.

MÉLANGES

Les mélanges extemporanés doivent être mis en œuvre conformément à la réglementation en vigueur et aux recommandations des guides de bonnes pratiques officiels.

SUPER BOUILLIE MACCLESFIELD 80 est compatible avec la plupart des produits phytopharmaceutiques couramment utilisés, à l'exception de ceux ayant une forte réaction acide ou alcaline. Pour les mélanges, il est donc recommandé d'effectuer un test préalable avec une petite quantité de produits avant de généraliser le mélange.

Protection de l'opérateur et du travailleur

Se laver les mains après toute manipulation/utilisation/intervention dans une parcelle préalablement traitée.

Ne pas manger, boire, téléphoner ou fumer lors de l'utilisation du produit.

Caractéristiques des EPI		PROTECTION DE L'UTILISATEUR PENDANT LES PHASES DE :			PROTECTION DU TRAVAILLEUR en cas d'intervention sur les parcelles traitées
		MÉLANGE/ CHARGEMENT et NETTOYAGE	APPLICATION AVEC PULVERISATEUR PORTÉ OU TRAINÉ À RAMPE, PNEUMATIQUE OU ATOMISEUR TRACTEUR AVEC OU SANS CABINE	APPLICATION MANUELLE LANCE OU PULVERISATEUR A DOS	
GANTS EN NITRILE réutilisables (certifiés EN 374-3) ou à usage unique (certifiés EN 374-2)		Réutilisables	À usage unique, si intervention sur le matériel de pulvérisation	Réutilisables	
EPI VESTIMENTAIRE 65% polyester / 35% coton ≥ 230 g/m² + traitement déperlant		EPI vestimentaire ET EPI partiel			
EPI PARTIEL blouse ou tablier à manches longues catégorie III type PB3 certifié EN14605+A1					
LUNETTES ou ECRAN FACIAL certifiés EN 166:2002 (CE, sigle 3)					
PROTECTION RESPIRATOIRE masque (EN 136:1998 ou EN 140:1998) équipé d'un filtre P3 (EN143:2006) minimum					
BOTTES certifiées EN 13 832-3:2006					

Pour protéger l'opérateur :

- pendant le mélange/chargement et le nettoyage du matériel de pulvérisation, porter :

- Gants en nitrile certifiés EN 374-3,
- Combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement déperlant.
- EPI partiel (blouse ou tablier à manches longues) de catégorie III et de type PB (3) à porter par-dessus la combinaison précitée,
- Lunettes ou écran facial certifiés norme EN 166 (CE, sigle 3),
- Protection respiratoire certifiée : conforme aux normes EN 136 (masque complet) ou EN 140 (demi-masque) et équipée d'un filtre P3 (EN 143) au minimum.

- pendant l'application, porter :

Application avec tracteur avec cabine :

- Combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement déperlant.
- Gants en nitrile certifiés EN 374-2 à usage unique dans le cas d'une intervention sur le matériel pendant la phase de pulvérisation. Dans ce cas, les gants ne doivent être portés qu'à l'extérieur de la cabine et doivent être stockés après utilisation à l'extérieur de la cabine.

Application avec tracteur sans cabine :

- Combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement déperlant.
- Gants en nitrile certifiés EN 374-2 à usage unique, dans le cas d'une intervention sur le matériel pendant la phase de pulvérisation.

Application manuelle avec une lance ou un pulvérisateur à dos :

- Combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement ~~déperlant~~.
- Gants en nitrile certifiés EN 374-3,
- Bottes de protection certifiées EN 13 832-3.

Pour protéger le travailleur :

Dans les cas où le travailleur serait amené à intervenir sur les parcelles traitées, porter une combinaison de travail en polyester 65% / coton 35% avec un grammage de 230 g/m² ou plus avec traitement ~~déperlant~~.

Rapporter les équipements de protection individuelle (EPI) usagés dans un sac translucide à votre distributeur partenaire ECO EPI ou faire appel à une entreprise habilitée pour la collecte et l'élimination de produits dangereux.

RÉSISTANCE

L'utilisation répétée, sur une même parcelle, de préparations à base de substances actives de la même famille chimique ou ayant le même mode d'action, peut conduire à l'apparition d'organismes résistants.

Pour réduire ce risque, l'utilisateur doit raisonner en premier lieu les pratiques agronomiques et respecter les conditions d'emploi du produit.

Il est conseillé d'alterner ou d'associer, sur une même parcelle, des préparations à base de substances actives de familles chimiques différentes ou à modes d'action différents, tant au cours d'une saison culturale que dans la rotation.

En dépit du respect de ces règles, on ne peut pas exclure une altération de l'efficacité de cette préparation liée à ces phénomènes de résistance. De ce fait, Industrias Químicas del Vallés décline toute responsabilité quant à d'éventuelles conséquences qui pourraient être dues à de telles résistances.

STOCKAGE

Conserver le produit uniquement dans son emballage d'origine, dans un local phytopharmaceutique conforme à la réglementation en vigueur, à l'écart des aliments et boissons, y compris ceux pour animaux.

Conserver hors de la portée des enfants et des personnes non autorisées.

NETTOYAGE DU PULVÉRISATEUR ET GESTION DES FONDS DE CUVE

À la fin de la période d'application avec le produit, l'intégralité de l'appareil (cuve, rampe, circuit, buses...) doit être rincée à l'eau claire additionnée d'un mouillant (recommandé pour le nettoyage des pulvérisateurs). Le rinçage du pulvérisateur, l'épandage ou la vidange du fond de cuve et l'élimination des effluents doivent être réalisés conformément à la réglementation en vigueur.

ÉLIMINATION DU PRODUIT ET DE SON EMBALLAGE

Réemploi de l'emballage interdit.

Apporter les emballages vidés et pliés à votre distributeur partenaire d'A.D.I.VALOR ou à un autre service de collecte spécifique.

Pour l'élimination des produits non utilisables, conserver le produit dans son emballage d'origine. Interroger votre distributeur partenaire d'A.D.I.VALOR ou faites appel à une entreprise habilitée pour la collecte et l'élimination des déchets dangereux.

En cas de déversement accidentel :

Se protéger (EPI) et sécuriser la zone. Prévenir les pompiers (18 ou 112) en cas de danger immédiat pour l'environnement que vous ne pouvez gérer avec vos propres moyens.

Collecter tout ce qui a pu être en contact avec le produit, terre souillée incluse. Nettoyer le site et le matériel utilisé, en prenant soin de confiner les effluents générés par l'opération de nettoyage. Les éliminer selon la réglementation en vigueur.



AVERTISSEMENT

Toute reproduction totale ou partielle de cette étiquette est interdite.

Respecter les usages, doses, conditions et précautions d'emploi mentionnés sur l'emballage. Ils ont été déterminés en fonction des caractéristiques du produit et des applications pour lesquelles il est préconisé. Conduire sur ces bases la culture et les traitements selon la bonne pratique agricole en tenant compte, sous la responsabilité de l'utilisateur, de tous les facteurs particuliers concernant votre exploitation, tels que la nature du sol, les conditions météorologiques, les méthodes culturales, les variétés végétales, la résistance des espèces...

Le fabricant garantit la qualité du produit vendu dans son emballage d'origine et stocké selon les conditions préconisées, ainsi que sa conformité à l'Autorisation de Mise sur le Marché délivrée par les autorités compétentes françaises. Pour les denrées issues de cultures protégées avec cette spécialité et destinées à l'exportation, il est de la responsabilité de l'exportateur de s'assurer de la conformité avec la réglementation en vigueur dans le pays importateur.