

REGISTRATION REPORT

Part A

Risk Management

Product code: A12738

Trade name : CALLIPRIME XTRA

Active Substance:

Mesotrione 480 g/L

COUNTRY: France

NATIONAL ASSESSMENT

Application for a label extension according to Art. 51

Minor uses

Applicant: Syngenta France

Date: 29/05/2017

Table des matières

1. DETAILS OF THE APPLICATION	3
1.1. Application background	3
1.2. Active substance approval	3
1.3. Regulatory Approach.....	3
1.4. Data Protection Claims.....	4
1.5. Letters of Access	4
2. DETAILS OF THE AUTHORISATION	4
2.1. Product Identity	4
2.2. Classification and Labelling	5
2.2.1. Classification and labelling in accordance with Regulation (EC) No 1272/2008	5
2.2.2. Other phrases in compliance with Regulation (EU) No 547/2011	5
2.2.3. Other phrases linked to the preparation.....	Erreur ! Signet non défini.
2.3. Product uses (Authorised GAP and uses in France).....	6
3. RISK MANAGEMENT	7
3.1. Reasoned statement of the overall conclusions taken in accordance with the Uniform Principles ..	7
3.1.1. Physical and chemical properties (Part B, Section 1, Points 2 and 4)	7
3.1.2. Methods of analysis.....	7
3.1.3. Mammalian toxicology.....	7
3.1.4. Residues and consumer exposure (Part B, Section 4)	7
Critical GAP(s) and overall conclusion.....	8
Summary of the evaluation.....	9
3.1.5. Efficacy (Part B, Section 7).....	10
3.2. Conclusions arising from French assessment.....	10
3.3. Substances of concern for national monitoring	10
3.4. Further information to permit a decision to be made or to support a review of the conditions and restrictions associated with the authorisation	10
Appendix 1 – Copy of the French decision	11
Appendix 2 – Copy of the draft product label as proposed by the applicant	16
Appendix 3 – Letter(s) of Access	17

PART A – Risk Management

The company SYNGENTA has requested a label extension according to article 51 in France for the product CALLIPRIME XTRA (A12738).

This document describes the specific conditions of use and labelling required for extension of the registration of CALLIPRIME XTRA (A12738) containing mesotrione.

The risk assessment conclusions are based on the already existing registration of the preparation in France. Therefore, the evaluation of the current application is limited to the points not covered by the existing registration.

Appendix 1 of this document provides a copy of the French decision.

Appendix 2 of this document is a copy of the draft product label as proposed by the applicant.

Appendix 3 of this document is a copy of the letter(s) of access.

1. DETAILS OF THE APPLICATION

1.1. Application background

CALLIPRIME XTRA (A12738) is a soluble concentrate (SC) containing 480 g a.s./L of mesotrione. It is used in agriculture as a herbicide. CALLIPRIME XTRA (A12738) is intended to be used as a herbicide on flax and linseed.

The complete GAP for the national application in France is provided below, under point 2.3.

1.2. Active substance approval

Mesotrione

Commission Implementing Regulation (EU) No 540/2011 of 25 May 2011 implementing Regulation (EC) No 1107/2009 of the European Parliament and of the Council as regards the list of approved active substances

Specific provisions of regulation were as follows :

Only use as herbicide may be authorised.

For the implementation of the uniform principles as referred to in Article 29(6) of Regulation (EC) No 1107/2009, the conclusions of the review report on mesotrione, and in particular Appendices I and II thereof, as finalised in the Standing Committee on the Food Chain and Animal Health on 15 April 2003 shall be taken into account.

An EFSA conclusion is available (2016;14(3):4419).

A Review Report is available (SANCO/1416/2001 -Final, 14 April 2003).

1.3. Regulatory Approach

The present application (n°2016-1452) was evaluated in France by the French Agency for Food, Environmental and Occupational Health & Safety (Anses)¹.

The current document based on Anses' assessment of the application submitted for this product is in compliance with Regulation (EC) no 1107/2009², implementing regulations and French regulations.

¹ French Food Safety Agency

² REGULATION (EC) No 1107/2009 of the European Parliament and of the Council of 21 October 2009 concerning the placing of plant protection products on the market and repealing Council Directives 79/117/EEC and 91/414/EEC

Since the application is intended for use in France only, the draft Part A was not circulated for comments. According to the French law and procedures, specific conditions of use are set out in the Decision letter. The French Order of 4 May 2017³ provides that:

- unless formally stated in the product authorisation, the pre harvest interval (PHI) is at least 3 days;
- unless formally stated in the product authorisation, the minimum buffer zone alongside a water body is 5 m;
- unless formally stated in the product authorisation, the minimum re-entry period is 6 hours for field uses and 8 hours for indoor uses.

Drift reduction measures such as low-drift nozzles are not considered within the decision making process in France. However, drift buffer zones may be reduced under some circumstances as explained in appendix 3 of the above-mentioned French order.

The data taken into account are those deemed to be valid either at European Union level or at zonal/national level. This part A presents a summary of essential scientific points upon which recommendations are based and is not intended to show the assessment in detail.

The conclusions relating to the acceptability of risk are based on the criteria indicated in Regulation (EU) N°546/2011⁴, and are expressed as “acceptable” or “not acceptable” in accordance with those criteria.

Finally, the French Order of 26 March 2014⁵ provides that:

- an authorisation granted for a « reference » crop applies also for “linked” crops unless formally stated in the decision
- the “reference” and “linked” crops are defined in appendix 1 of that French order.

Thus, at French national level, possible extrapolation of submitted data and the corresponding assessment from “reference” crops to “linked” ones are undertaken even if not clearly requested by the applicant in their dRR, and a conclusion is reached on the acceptability of the intended uses on those “linked” crops. The aim of this Order, mainly based on the EU document on residue data extrapolation⁶ is to supply “minor” crops with registered plant protection products.

Therefore the GAP table (Section 2.3) and Decision may include uses on crops not originally requested by the applicant. The Decision, as reproduced in Appendix 1, takes also into account national provisions, including national mitigation measures.

1.4. Data Protection Claims

There is no new data submitted with this application.

1.5. Letters of Access

Not relevant for this application.

2. DETAILS OF THE AUTHORISATION

2.1. Product Identity

³

<https://www.legifrance.gouv.fr/eli/arrete/2017/5/4/AGR1632554A/jo/texte>

⁴ COMMISSION REGULATION (EU) No 546/2011 of 10 June 2011 implementing Regulation (EC) No 1107/2009 of the European Parliament and of the Council as regards uniform principles for evaluation and authorisation of plant protection products

⁵ <http://www.legifrance.gouv.fr/eli/arrete/2014/3/26/AGR1407093A/jo>

⁶ SANCO document “guidance document:- Guidelines on comparability, extrapolation, group tolerances and data requirements for setting MRLs”: SANCO/ 7525/VI/95 - rev.9

Product Name	CALLIPRIME XTRA (A12738)
Authorization Number (for re-registration)	2150012
Function	Herbicide
Applicant	SYNGENTA FRANCE SAS
Composition	480 g/L of mesotrione
Formulation type	Soluble Concentrate [Code: SC]
Packaging	Not relevant for extension of authorisation according article 51.

2.2. Classification and Labelling

2.2.1. Classification and labelling in accordance with Regulation (EC) No 1272/2008

Not relevant for extension of authorisation according article 51.

Additional labelling phrases: Contains 1,2-benzisothiazolin-3(2H)-one

2.2.2. Other phrases in compliance with Regulation (EU) No 547/2011

This phrase should be added:

SPe 3	To protect aquatic organisms respect an unsprayed vegetated buffer zone of 20 meters to surface water bodies.
SPe 3	To protect non-target plants respect an unsprayed buffer zone of 20 meters to non-agricultural land.

Please refer to the decision of product authorization.

2.2.3. Other phrases linked to the preparation

Wear suitable personal protective equipment ⁷ : refer to the Decision in Appendix 1 for the details.
Re-entry period : refer to the decision of product authorisation.
Pre-harvest interval :
- linseed : F – last application on BBCH 05.
Other mitigation measures: refer to the decision of product authorisation.
The label must reflect the conditions of authorisation.

⁷ If a tractor with cab is used, wearing gloves during application is only required when working with the spray mixture

2.3. Product uses (Authorised GAP and uses in France)

PPP (product name/code)	CALLIPRIME XTRA (A12738) mesotrione	Formulation type: SC
Applicant:	Syngenta	Conc. of as 1: <input checked="" type="checkbox"/> professional use <input type="checkbox"/> non professional use
Zone(s):	south EU	<input checked="" type="checkbox"/> <input type="checkbox"/>

Verified by MS: FR

1	2	3	4	5	6	7	8	9	10	11	12	13
Use- No.	Member state(s)	Crop and/ or situation (crop destination / purpose of crop)	F G or I	Pests or Group of pests controlled (additionally: developmental stages of the pest or pest group)	Application			Application rate			PHI (days)	Remarks: e.g. safener/synergist per ha e.g. recommended or mandatory tank mixtures
					Method / Kind	Timing / Growth stage of crop & season	Max. number (min. interval between applications) a) per use b) per crop/ season	kg. L product / ha	g, kg as/ha	Water L/ha min / max		
1	FR	Flax	F	Grass and dicotyledoneous weed	Foliar spray	BBCH 00 - 05	1	0,31 l/ha	150	100 - 400		Acceptable.
2	FR	Linseed,	F	Grass and dicotyledoneous weed	Foliar spray	BBCH 00 - 05	1	0,31 l/ha	150	100 - 400		Acceptable.

Remarks table:

- (1) Numeration necessary to allow references
- (2) Use official codes/nomenclatures of EU
- (3) For crops, the EU and Codex classifications (both) should be used; where relevant, the use situation should be described (e.g. fumigation of a structure)
- (4) F: professional field use, Fn: non-professional field use, Fpn: professional and non-professional field use, G: professional greenhouse use, Gn: non-professional greenhouse use, Gpn: professional and non-professional greenhouse use,
- (5) Scientific names and EPPO-Codes of target pests/diseases/ weeds or when relevant the common names of the pest groups (e.g. biting and sucking insects, soil born insects, foliar fungi, weeds) and the developmental stages of the pests and pest groups at the moment of application must be named
- (6) Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench
Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plants - type of equipment used must be indicated
- (7) Growth stage at first and last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application
- (8) The minimum and maximum number of application possible under practical conditions of use must be provided
- (9) For specific uses other specifications might be possible, e.g.: g/m³ in case of fumigation of empty rooms. See also EPPO-Guideline PP 1/239 Dose expression for plant protection products
- (10) The dimension (g, kg) must be clearly specified. (Maximum) dose of a.s. per treatment (usually g, kg or L product / ha).
- (11) If water volume range depends on application equipments (e.g. ULVA or LVA) it should be mentioned under “application: method/kind”.
- (12) PHI - minimum pre-harvest interval
- (13) Remarks may include: Extent of use/economic importance/restrictions

3. RISK MANAGEMENT

3.1. Reasoned statement of the overall conclusions taken in accordance with the Uniform Principles

3.1.1. Physical and chemical properties (Part B, Section 1, Points 2 and 4)

Not relevant for extension of authorisation according article 51.

3.1.2. Methods of analysis

3.1.2.1 Analytical method for the formulation

Not relevant for extension of authorisation according article 51.

3.1.2.2 Analytical methods for residues

Further data for this application are not necessary.

3.1.3. Mammalian toxicology

Endpoints used in risk assessment

Active Substance: Mesotrione			
ADI	0.01 mg / kg bw/d		
ARfD	0.02 mg/kg bw		EU (2003)
AOEL	0.015 mg/kg bw/d		
AOEL	0.005 mg/kg bw/d		EFSA (2016)
Dermal absorption	Based on values adopted during the product authorization.		
		Concentrate	Spray dilution
	Dermal absorption endpoints %	0.2	2

3.1.3.1 Acute toxicity

CALLIPRIME XTRA (A12738) containing 480 g/L mesotrione has a low toxicity in respect to acute oral, inhalation and dermal toxicity and is not irritating to the rabbit skin or eye and is a skin sensitizer.

Nature of protective clothing and PPE for the operator:

Refer to the French decision.

Nature of protective clothing and PPE for the worker:

Refer to the French decision.

Re-entry period: 48 hours

3.1.3.2 Operator Exposure

Summary of critical use patterns (worst cases):

Crop	F/G ⁸	Equipment	Application rate kg/L product/ha (g as/ha)	Spray dilution (L/ha)	Model
Linen	F	Tractor-mounted boom sprayer-hydraulic nozzle	0.33 L/ha (158 g as/ha)	80	BBA

Considering intended use, operator systemic exposure was estimated using the German BBA model

Crop	Equipment	PPE and/or working coverall	% AOEL Mesotrione 0.015 mg/kg bw/d	% AOEL Mesotrione 0.005 mg/kg bw/d
Linen	Tractor-mounted boom sprayer-hydraulic nozzle	Working coverall and gloves during mixing/loading and application	2.2	6.7

Operator exposure was already assessed for similar uses in open field at higher application rate. It is concluded that the risk for the operator using CALLIPRIME XTRA (A12738) is acceptable with a working coverall (90% protection factor) and gloves during mixing/loading and application.

3.1.3.3 Bystander Exposure

Bystander exposure has been previously assessed according to the EUROPOEM II. It is concluded that there is no unacceptable risk for the bystander after incidental short-term exposure to CALLIPRIME XTRA (A12738).

Former AOEL:

Bystander exposure was assessed according to EUROPOEM II. Exposure is estimated to 0.7 % of the AOEL of mesotrione.

New AOEL

Bystander exposure was assessed according to EUROPOEM II. Exposure is estimated to 2.1 % of the AOEL of mesotrione.

It is concluded that there is no unacceptable risk to the bystander after incidental short-term exposure to CALLIPRIME XTRA (A12738).

3.1.3.4 Worker Exposure

CALLIPRIME XTRA (A12738) is used as herbicidal treatment on several crops where there is no need to re-enter the treated area after application. Worker exposure is considered not relevant.

3.1.4. Residues and consumer exposure (Part B, Section 4)

Critical GAP(s) and overall conclusion

Overall conclusion

The data available are considered sufficient for risk assessment. An exceedance of the current MRL for mesotrione as laid down in Reg. (EU) 396/2005 is not expected.

⁸ Open field or glasshouse

The chronic and the short-term intakes of mesotrione residues resulting from the uses intended in the framework of this application are unlikely to present a public health concern.

As far as consumer health protection is concerned, France agrees with the authorization of the intended use.

According to available data, no specific mitigation measures should apply.

Data gaps

None

Data required in post-authorization

None

Summary of the evaluation

The preparation CALLIPRIME XTRA (A12738A) is composed of mesotrione.

Summary for mesotrione

Use-No.*	Crop	Plant metabolism covered?	Sufficient residue trials?	PHI sufficiently supported?	Sample storage covered by stability data?	MRL compliance Reg (EU) No 2016/53	Chronic risk for consumers identified?	Acute risk for consumers identified?	Comments
1	Flax	Yes	NR	NR	NR	NR	No	NR	Non-alimentary use
2	Spring linseed	Yes	Yes (4 NEU)	Yes	Yes				/

NR: Not Relevant

* Use number(s) in accordance with the list of all intended GAPs in Part B, Section 0 should be given in column 1

As residues of linseed do not exceed the trigger values defined in Reg (EU) No 283/2013, there is no need to investigate the effect of industrial and/or household processing.

Residues in succeeding crops have been sufficiently investigated taking into account the specific circumstances of the cGAP being considered here. It is very unlikely that residues will be present in succeeding crops.

Considering dietary burden and based on the intended uses, no significant modification of the intake was calculated for livestock. Further investigation of residues as well as the modification of MRLs in commodities of animal origin is therefore not necessary.

Information on CALLIPRIME XTRA (A12738A)

Crop	PHI for CALLIPRIME XTRA (A12738A) proposed by applicant	PHI/ Withholding period* sufficiently supported for	PHI for CALLIPRIME XTRA (A12738A) proposed by zRMS	zRMS Comments (if different PHI proposed)
		Mesotrione		
Flax & Linseed	F**	Yes	F**	/

* Purpose of withholding period to be specified

** F: PHI is defined by the application stage at last treatment (time elapsing between last treatment and harvest of the crop).

Crop	PHI for CALLIPRIME XTRA (A12738A) proposed by applicant	PHI/ Withholding period* sufficiently supported for	PHI for CALLIPRIME XTRA (A12738A) proposed by zRMS	zRMS Comments (if different PHI proposed)
		Mesotrione		

Waiting periods before planting succeeding crops

Not relevant

1.1.1 Environmental fate and behaviour

The fate and behaviour in the environment have been evaluated according to the requirements of Regulation (EC) No 1107/2009. Appropriate endpoints from the EU conclusions were used to calculate PEC values for the active substance and its metabolites for the intended use patterns. In cases where deviations from the EU agreed endpoints were considered appropriate (for example when additional studies are provided), such deviations were highlighted and justified accordingly.

No PECsoil, PECgw or PECsw are available for mesotrione and its metabolites specifically for the use on linseed. The risk of groundwater contamination cannot be finalised for this use.

Based on vapour pressure, information on volatilisation from plants and soil, and DT50 calculation, no significant contamination of the air compartment is expected for the intended uses.

1.1.2 Ecotoxicology

The ecotoxicological risk assessment of the formulation was performed according to the requirements of Regulation (EC) No 1107/2009. Appropriate endpoints from the EU conclusions for the active substance(s) and its/their metabolites were used for the intended use patterns. In cases where deviations from the EU agreed endpoints were considered appropriate (for example when additional studies are provided), such deviations were highlighted and justified accordingly.

Based on the guidance documents, the risks for birds, bees and other non-target arthropods, earthworms, other soil macro-organisms and micro-organisms and terrestrial plants are acceptable for the intended uses. Since no PECsw are available in E-fate Section, the risk for aquatic organisms cannot be finalized. For mammals, the refined risk assessment based on specific data for maize from the original evaluation of this formulated product cannot be used for the current intended uses on linseed. Therefore, the risk for mammals cannot be finalized.

3.1.5. Efficacy (Part B, Section 7)

According to Article 51 of Regulation (EC) No 1107/2009, the efficacy assessment and the absence of any phytotoxicity risk on the crop is not necessary.

3.2. Conclusions arising from French assessment

Taking into account the above assessment, even if the risk of groundwater contamination, for aquatic organisms and mammals was not finalised, considering that flax and linseed are minor crops in France, an authorisation was granted. A copy of the decision issued can be found in Appendix 1 – Copy of the product decision.

3.3. Substances of concern for national monitoring

No information stated.

3.4. Further information to permit a decision to be made or to support a review of the conditions and restrictions associated with the authorisation

No further information is required.

Appendix 1 – Copy of the French decision



Décision relative à une demande d'extension d'usage(s) d'un produit phytopharmaceutique

Vu les dispositions du règlement (CE) N° 1107/2009 du 21 octobre 2009 et de ses textes d'application,

Vu le code rural et de la pêche maritime, notamment le chapitre III du titre V du livre II des parties législative et réglementaire,

Vu la demande d'extension d'usage mineur du produit phytopharmaceutique CALLIPRIME XTRA

de la société SYNGENTA FRANCE S.A.S.
enregistrée sous le n°2016-2682

Vu les conclusions de l'évaluation de l'Anses du 28 avril 2017,

L'autorisation de mise sur le marché du produit référencé ci-après **est étendue** aux usages décrits dans la présente décision.

La présente décision s'applique sans préjudice des autres dispositions applicables.

Avertissement :

Le non-respect des conditions décrites ci-dessous peut entraîner le retrait ou la modification de l'autorisation ainsi que toute action incluant des poursuites judiciaires.



Informations générales sur le produit


Noms du produit	CALLIPRIME XTRA CALLISTO 480 SC LUMESTRA 480
Type de produit	Produit de référence
Titulaire	SYNGENTA FRANCE S.A.S. 12 Chemin de l'Habit 31790 Saint Sauveur FRANCE
Formulation	Suspension concentrée (SC)
Contenant	480 g/L - mésotrione
Numéro d'intrant	2150033
Numéro d'AMM	2150012
Fonction	Herbicide
Gamme d'usages	Professionnel

L'échéance de validité de la présente décision correspond à celle de l'autorisation du produit.

La présente décision peut être retirée ou modifiée si des éléments le justifient.

A Maisons-Alfort, le 29 MAI 2017

Françoise WEBER
Directrice générale adjointe des produits réglementés
Agence nationale de sécurité sanitaire de
l'alimentation, de l'environnement et du travail (ANSES)



ANNEXE I : Modalités d'autorisation du produit

Liste des usages autorisés

En l'absence de restriction, les usages sont autorisés sur l'ensemble des cultures de la portée de l'usage.

Usages	Dose maximale d'emploi	Nombre maximum d'applications	Stade d'application BBCH	Délai avant récolte (jours)	Zone Non Traitée aquatique (mètres)	Zone Non Traitée arthropodes non cibles (mètres)	Zone Non Traitées plantes non cibles (mètres)	Mention abeilles
15505902 Lin-Désherbage	0,31 L/ha	1/an	entre les stades BBCH 00 et BBCH 05	F (BBCH 05)	20 (dont DVP 20)	-	20	-

Autorisé sur lin textile et oléagineux.

DVP : Dispositif Végétalisé Permanent.

CALLIPRIME XTRA
AMM n°2150012



Conditions d'emploi du produit

Protection de l'opérateur et du travailleur

Pour l'opérateur, porter

Dans le cadre d'une application avec pulvérisateur à rampe :

• pendant le mélange/chargement

- Gants en nitrile certifiés EN 374-3 ;
- Combinaison de travail tissée en polyester 65 %/coton 35 % avec un grammage de 230 g/m² ou plus avec traitement déperlant ;
- EPI partiel (blouse) de catégorie III et de type PB (3) à porter par-dessus la combinaison précitée

• pendant l'application

Si application avec tracteur avec cabine

- Combinaison de travail tissée en polyester 65 %/coton 35 % avec un grammage de 230 g/m² ou plus avec traitement déperlant ;
- Gants en nitrile certifiés EN 374-3 à usage unique, dans le cas d'une intervention sur le matériel pendant la phase de pulvérisation. Dans ce cas, les gants ne doivent être portés qu'à l'extérieur de la cabine et doivent être stockés après utilisation à l'extérieur de la cabine.

Si application avec tracteur sans cabine

- Combinaison de travail tissée en polyester 65 %/coton 35 % avec un grammage de 230 g/m² ou plus avec traitement déperlant ;
- Gants en nitrile certifiés EN 374-3 à usage unique pendant l'application et dans le cas d'une intervention sur le matériel pendant la phase de pulvérisation ;

• pendant le nettoyage du matériel de pulvérisation

- Gants en nitrile certifiés EN 374-3 ;
- Combinaison de travail tissée en polyester 65 %/coton 35 % avec un grammage de 230 g/m² ou plus avec traitement déperlant ;
- EPI partiel (blouse) de catégorie III et de type PB (3) à porter par-dessus la combinaison précitée ;

Pour le travailleur, amené à entrer dans la culture après traitement, porter

- Une combinaison de travail (cotte en coton/polyester 35 %/65 % - grammage d'au moins 230 g/m²) avec traitement déperlant.

Délai de rentrée

- 48 heures en application de l'arrêté du 4 mai 2017.

Respect des limites maximales de résidus (LMR)

- Pour chaque usage figurant dans la liste des usages autorisés, les conditions d'utilisation du produit permettent de respecter les limites maximales de résidus.

Protection de l'environnement (milieux, faune et flore)

Protection de la faune

Pour protéger les organismes aquatiques, respecter une zone non traitée de 20 mètres en bordure des points d'eau comportant un dispositif végétalisé permanent non traité d'une largeur de 20 mètres.



Protection de la flore

Pour protéger les plantes non cibles, respecter une zone non traitée de 20 mètres par rapport à la zone non cultivée adjacente.

Les autres conditions d'emploi du produit restent inchangées.

Recommandations relatives à l'étiquette du produit

Contient de la 1,2-benzisothiazolin-3(2H)-one.

Appendix 2 – Copy of the draft product label as proposed by the applicant

CALLIPRIME XTRA (AMM n° 2150012) Extension d'usage sur lin – Projet d'étiquette

Les tableaux ci-dessous viennent se substituer aux tableaux de l'étiquette existante.

Etiquette DOS ou Pavé réglementaire :

CULTURES AUTORISEES, UNIQUEMENT :	CIBLES	DOSES AUTORISEES	DELAI AVANT RECOLTE
Maïs	Désherbage	0,33 L/ha	Traiter jusqu'au stade BBCH 14 au plus tard
Maïs doux	Désherbage	0,33 L/ha	42 jours
Sorgho	Désherbage	0,2 L/ha	Traiter jusqu'au stade BBCH 14 au plus tard
Lin textile et lin oléagineux de printemps	Désherbage	0,31 L/ha	Traiter jusqu'au stade BBCH 05 du lin oléagineux au plus tard

Etiquette Livret (page 3 du PDF joint) :

CULTURES AUTORISEES, UNIQUEMENT :	CIBLES	DOSES AUTORISEES	NOMBRES MAX. D'APPLICATIONS	Délai Avant Récolte ou Période d'application	ZNT ^{**}	DVP ^{***}
Maïs (grains et fourrage) Maïs semences*	Désherbage	0,33 L/ha	1	Du stade pré- semis jusqu'au stade 4 feuilles vraies du maïs (BBCH 14 maxi) 42 jours Du stade post- semis / prélevée, jusqu'au stade 4 feuilles vraies du maïs doux	20 mètres par rapport à la zone non cultivée adjacente et 20 mètres en bordure des points d'eau temporaires ou permanents	20 mètres en bordure des points d'eau temporaires ou permanents
Maïs doux	Désherbage	0,33 L/ha	1	post-semis / prélevée, jusqu'au stade 4 feuilles vraies du sorgho (BBCH 14 maxi)		
Sorgho	Désherbage	0,2 L/ha	1	Jusqu'au stade BBCH 05 du lin oléagineux		
Lin textile et lin oléagineux de printemps	Désherbage	0,31 L/ha	1			

* Avec accord de l'horticulteur

** Zone Non Traitée

*** Dispositif Végétalisé Permanent non traité

Appendix 3 – Letter(s) of Access

Not applicable.